

FARMAG

CYPERMETHRIN

Reg. No. / Nr L3207 Act No. 36 of 1947/ Wet Nr 36 van 1947



INSECTICIDE

**READ THE LABEL AND ATTACHED PACKAGED LEAFLET BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

IRAC INSECTICIDE GROUP CODE

3

IRAC INSEKDODER GROEPKODE

An emulsifiable concentrate contact and stomach insecticide for the control of agricultural pests as indicated.

'n Emulgeerbare konsentraat kontak- en maaginsekodder vir die beheer an landbouplae soos aangedui.

ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL:

Cypermethrin (pyrethroid)200 g/l..... Sipermetrien (piretroïed)



DANGER

Batch No:

D.O.M:

D.O.E:

Net Mass:

See reverse of packaging for details.

FOR FULL PARTICULARS SEE ATTACHED PAMPHLET.
VIR VOLLEDIGE BESONDERHEDE KYK AANGEHEGTE ETIKET.

UN No. / VR Nr: 3351



REGISTRATION HOLDER/REGISTRASIEHOOUER
Castle Ag-Chem (Pty) Ltd.
Reg. No. / Nr. 2007/033120/07
61 Marshall Drive, Old Mill Industrial Park,
Mount Edgecombe.
Tel: (031) 003 3486

Hazard Statements:

Flammable liquid and vapour.
Toxic if swallowed.
Harmful in contact with skin or if inhaled.
Causes skin irritation.
May cause respiratory irritation.
May cause damage to organs through prolonged or repeated exposure.
Very toxic to aquatic life.
Toxic to aquatic life with long lasting effects.

Precautionary Statements:

Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking.
Do not breathe vapours, spray.
IF SWALLOWED: Immediately call a poison center or doctor.
IF ON SKIN: Wash with plenty of water.
IF INHALED: Remove person to fresh air and keep comfortable for breathing.
If skin irritation occurs: Get medical advice/attention.
In case of fire: Use carbon dioxide (CO₂), dry extinguishing powder, foam to extinguish.

Poison Information Helpline (24 hrs): 0861 555 777



BARCODE

100450

165(H) x 140(W) mm

CYAN MAG YEL BLK

100450

WARNINGS

Hazard Statements: Flammable liquid and vapour. Toxic if swallowed. Harmful in contact with skin or if inhaled. Causes skin irritation. May cause respiratory irritation. May cause damage to organs through prolonged or repeated exposure. Very toxic to aquatic life. Toxic to aquatic life with long lasting effects

PRE-HARVEST INTERVAL:

Withholding periods: minimum number of days between last application and harvest or grazing:

Apples and Pears.....	14 days
Beans.....	7 days
Cotton (grazing).....	28 days
Cruciferae.....	4 days
Grapes (wine and table).....	28 days
Groundnuts (hay).....	7 days
Lucerne (grazing).....	14 days
Macadamia nuts	30 days
Maize.....	14 days
Pastures (grazing).....	14 days
Peaches.....	14 days
Peas.....	7 days
Sorghum.....	28 days
Sweetcorn.....	14 days
Tomatoes.....	4 days
Citrus.....	28 days

- **Re-entry Interval:** Do not enter treated area within 1 day after treatment, unless wearing protective clothing.
- **In case of poisoning call a doctor and make the label available to him.**
- **Aerial application warning:** Notify all inhabitants of the immediate area to be sprayed and issue the necessary warning. Do not spray over or allow drift to contaminate adjacent areas, rivers or dams.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, incompatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS

Precautionary Statements

General: - If medical advice is needed, have product container or label at hand. Keep out of reach of children. Read label before use.

Prevention: Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking. Keep container tightly closed. Ground and bond container and receiving equipment. Use explosion-proof equipment. Use non-sparking tools. Take action to prevent static discharges. Do not breathe vapours, spray. Wash hands, forearms and face thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection.

Response: IF SWALLOWED: Immediately call a poison center or doctor. IF ON SKIN: Wash with plenty of water. Take off immediately all contaminated clothing. Rinse skin with water/shower. IF INHALED: Remove person to fresh air and keep comfortable for breathing. Call a poison center or doctor if you feel unwell. Get medical advice/attention if you feel unwell. Specific treatment (see supplemental first aid instruction on this label). Rinse mouth. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse. In case of fire: Use carbon dioxide (CO₂), dry extinguishing powder, foam to extinguish. Collect spillage

Storage: Keep container tightly closed. Store in a well-ventilated place. Keep cool. Store locked up.

Disposal: Dispose of contents and container to an approved waste disposal plant.

- Avoid drift of spray mist onto other crops, grazing, rivers and dams and areas not under treatment.
- Clean application equipment after use. Dispose of wash water where it will not contaminate crops, grazing, rivers, dams and boreholes.
- **TRIPLE RINSE the empty container as follows:** Invert the empty container over the spray or mixing tank and drain for at least 30 seconds after the flow has slowed down to dripping. Thereafter rinse the empty container three times in succession with one quarter of the container volume fresh water and decant the rinsate into the spray or mixing tank. Puncture the triple rinsed container and dispose of via an approved collector or recycler (www.croplife.co.za). Do not bury, burn or donate the container to any other parties that may use it as a container for food or beverages.
- Destroy empty container by perforation, flattening and burying.
- Do not use empty container for any other purpose.
- Prevent contamination of food, feed, eating utensils and drinking water.

RELEVANT SUBSTANCES:

NAME	PRODUCT IDENTIFIER
Cypermethrin 97% TC	CAS-No.: 52315-07-8
Xylene	CAS-No.: 1330-20-7
Tristyrylphenol Ethoxylates	CAS-No.: 99734-09-5
Calcium dodecylbenzenesulphonate	CAS-No.: 26264-06-2

SYMPOTMS OF POISONING

Causes burning, itching or tingling sensations of the skin which readily disappears within 24 hours after exposure. Inhalation causes nasal discharge and a scratchy throat. Symptoms of high-dose ingestion include nausea, prolonged vomiting, stomach pains and diarrhea progressing to convulsions, unconsciousness and coma. Hypersensitivity, tremors, muscular fibrillation, salivation, in co-ordination, incontinence of urine, and lacrimation, rapid and /or irregular respiration.

FIRST AID TREATMENT

Eyes: Flush eyes with clean, gentle flowing water for at least 20 minutes, occasionally lifting upper and lower lids, until no evidence of chemical remains. If irritation occurs and persists, obtain medical attention.

Skin: Remove contaminated clothing and rinse contaminated body area thoroughly with plenty of soap and cold water, until no evidence of chemical remains. Do not rub skin. Apply olive oil to affected areas for prompt relief. Obtain medical attention if irritations occurs or persists.

Inhalation: Move the patient to fresh air. If breathing is difficult or irritation occurs and persists, obtain medical attention.

Ingestion: Do not induce vomiting. If patient is alert, rinse mouth thoroughly with clean water. Do not give anything by mouth if patient is unconscious. If vomiting occurs, prevent vomit being inhaled. Obtain medical attention.

NOTE TO PHYSICIAN

This pesticide contains a pyrethroid and an aromatic solvent. If pesticide is aspirated into lungs during ingestion or vomiting, mild to severe chemical pneumonia may be caused. The stomach should be emptied as soon as possible by careful gastric lavage.

No antidote is available. Treat symptomatically and supportively. In case of ingestion, consider gastric lavage with water or 5 % sodium bicarbonate solution. Convulsions should be treated with anti-convulsants.

RESISTANCE WARNING

For resistance management, **FARMAG CYPERMETHRIN** is a group code 3 insecticide. Any insect population may contain individuals naturally resistant to **FARMAG CYPERMETHRIN** and other group code 3 insecticides. The resistant individuals can eventually dominate the insect population if these insecticides are used repeatedly. These resistant insects may not be controlled by **FARMAG CYPERMETHRIN** or any other group code 3 insecticides.

To delay insecticide resistance:

- Avoid exclusive repeated use of insecticide from the same insecticide group code. Alternate or tank mix with products from different insecticide group codes.
- Integrate other control methods (chemical, cultural, biological) into insect control programmes.

For specific information on resistance management contact the registration holder of this product.

Warning against American (African) bollworm resistance:

Resistance of American (Africa) bollworm (*Helicoverpa armigera*) to synthetic pyrethroids has been confirmed.

Cotton: Synthetic pyrethroids must only be applied to cotton during the period 1 January to 28 February.

All other crops: Do not apply more than two applications per growing season. If a pyrethroid gives poor performance, do not respray with any synthetic pyrethroid, even at a corrective dosage rate.

Use a product from a different chemical group.

MODE OF ACTION:

Acts on the nervous system of the insect, and disturbs the function of neurons by interaction with the sodium channel. Non-systemic insecticide with contact and stomach action. Also exhibits anti-feeding action. Good residual activity on treated plants.

DIRECTIONS FOR USE: Use only as indicated.**Compatibility**

- Do not mix **FARMAG CYPERMETHRIN** with seaweed extract, vamidothion, prothiophos, chinomethionate and amitraz containing pesticides.
- Do not add a wetting agent to the **FARMAG CYPERMETHRIN** spray mixture when spraying apples, pears and grapes.
- **FARMAG CYPERMETHRIN** is compatible with a wide range of pesticides. However, since it is not possible for Cipla Agricare (Pty) Ltd to test all possible combinations, it is advised to carry out a compatibility test on a small scale in the event of any uncertainty.
- The effectiveness of **FARMAG CYPERMETHRIN** can be negatively influenced by high pH values of the spray mixture. Cypermethrin, the active ingredient of **FARMAG CYPERMETHRIN**, is more stable at a pH of around 4. Where an acidifying agent is used to lower the pH of the water, the agent must be added to the water before adding **FARMAG CYPERMETHRIN**.
- The addition of molasses to the spray mixture has an acidifying effect and is used to reduce evaporation of the spray mist. For ground application add 10 % molasses by volume and 20 % for aerial application.

Mixing Instructions

Half fill the spray tank with clean water. Add the required amount of **FARMAG CYPERMETHRIN** to the tank whilst agitating. Thereafter, fill the tank to the required volume with water and maintain agitation until spraying is completed. Refer 'COMPATIBILITY' above regarding pH of spray mixture.

Ground Application

When using a conventional high volume sprayer or a mistblower, fit equipment with hollow cone nozzles that give medium to fine droplets. Ensure that sprayers are calibrated before spraying and care taken of thorough coverage of the crop and uniform distribution of spray mixture.

Volume water to be applied per hectare in deciduous fruit orchards:

The volume of water to be applied per hectare in deciduous fruit orchards is calculated according to the Tree Row Volume Formula as described by Unrath:

The formula is as follows: $\text{Tree height} \times \text{tree diameter} \times 937 = \text{Volume water to be applied per hectare}$

Row width

The measurements used in the formula are those of the tree in full leaf cover. Because deciduous fruit trees are not always in full leaf the water volume to be applied per hectare at the earlier growth stages is as follows:

From green-tip to 30 % blossom - 60 % of HV to be applied From ca. full blossom to ca.

November - 80 % of HV to be applied

From ca. beginning of December - 100 % of HV to be applied

When pesticides are applied as low volume sprays the correct quantity of pesticide, as calculated for high volume applications, must be applied per hectare.

Aerial Application

Aerial application of **FARMAG CYPERMETHRIN** may only be done by a registered Aerial Application Operator using a correctly calibrated, registered aircraft according to the instructions of SANS 10118: 2009 (The Aerial Application of Pesticides). Ensure that the spray mixture is distributed evenly over the target area and that the loss of spray material during application is restricted to a minimum. It is therefore essential that the following criteria be met:

- **Volume:** A spray mixture volume of 30 litre per hectare is recommended. As this product has not been evaluated at a reduced volume rate, the registration holder cannot guarantee efficacy, or be held responsible for any adverse effects if this product is applied aerially at a lower volume rate than recommended above.

- **Droplet coverage:** 30 to 40 droplets per cm² must be recovered at the target area.
- **Droplet size:** A droplet spectrum with a VMD of 250 to 280 microns is recommended. Limit the production of fine droplets less than 150 microns (high drift and evaporation potential) to a minimum.
- **Flying height:** Maintain the height of the spray boom at 3 to 4 metres above the target. Do not spray when aircraft dives, is in a climb or when banking.
- Use suitable **atomising equipment** that will produce the desired droplet size and coverage, but which will ensure the minimum loss of product. The spraying system must produce a droplet spectrum with the lowest possible Relative Span.
- Position all the atomisers within the inner 60 to 75 % of the wingspan to prevent droplets from entering the **wingtip vortices**.
- The difference in **temperature** between the wet and dry bulb thermometers, of a whirling hygrometer, should not exceed 8 °C.
- Stop spraying if the **wind speed** exceeds 15 km/h.
- Stop spraying under **turbulent**, unstable and dry conditions during the heat of the day.
- Spraying under temperature **inversion conditions** (spraying in or above the inversion layer) and/or **high humidity conditions** (relative humidity 80 % and above) may lead to the following:
 - reduced efficacy due to suspension and evaporation of small droplets in the air (inadequate coverage).
 - damage to other sensitive crops and/or non-target areas through drifting of the suspended spray cloud away from the target field.
- Ensure that the Aerial Spray Operator knows exactly which fields to spray.
- Obtain an assurance from the Aerial Spray Operator that the above requirements will be met and that relevant data will be compiled in a logbook and kept for future reference.

CROP/PEST	DOSAGE RATE / 100L WATER OR AS INDICATED	REMARKS
AFORRESTATION Pine emperor moth larva	50 mL/ha	Aerial application Apply in 30L water per ha
Willow tree emperor moth larva	100 mL/ha	Aerial application Apply in 30L water per ha
Wattle bagworm	100 mL/ha	Aerial application Apply in 30L water per ha commencing November/December
ALL CROPS Cutworms	100 mL/ha 0.33 mL/100m plant row 100 mL/ha	Apply as soon as pest is noticed as a treatment either pre- or post-emergence of crop. Ensure that top 3cm of soil is moist. Repeat treatment in cases of re-infestation. Ground Application Overall spray: apply in 300L water per ha. Band spray (300 mm breed): apply in at least 3L water per 100 m row. Aerial application Apply in 30L water per ha.
APPLES & PEARS		To determine the volume of water in which FARMAG CYPERMETHRIN 200 EC should be applied per hectare refer ' GROUND APPLICATION: Volume water to be applied per hectare in deciduous fruit orchards '. Also refer ' COMPATIBILITY '.
Codling moth Leaf roller	5 mL	Apply first spray at 75% petal drop and repeat at 14 day intervals.
American (Africa) bollworm	5 mL	Apply as a corrective treatment not earlier than 75% petal drop. Refer ' Warning against American (Africa) bollworm resistance ' under ' RESISTANCE WARNING '
Banded fruit weevil (Snout beetle)	10 mL	Apply two sprays, the first 75% petal drop. Repeat 4 weeks later. Further, either a low or high volume spray can be applied during mid-January to prevent late season damage.

BEANS American (Africa) bollworm	150 mL/ha	<p>Ground application Apply in 300L water per ha. Inspect lands regularly and if necessary, repeat application.</p> <p>Aerial application Apply in 30L water per ha. Refer 'Warning against American (Africa) bollworm resistance' under 'RESISTANCE WARNING'</p>
COTTON American (Africa), red and spiny bollworms Stainers Semi- (Plusia) looper	150 mL/ha	<p>See 'Warning against American (Africa) bollworm resistance' under 'RESISTANCE WARNING'. In order to comply with the principles of pest management and integrated control measures, FARMAG CYPERMETHRIN is intended to be used during the period of fruiting from peak flowering until boll split, i.e. approximately 10 to 22 weeks after crop emergence. FARMAG CYPERMETHRIN is primarily intended for use as a preventive control programme applied at 7 day intervals. Normally a bollworm spray programme will commence at the beginning of flowering i.e. about 6 weeks after crop emergence. From then until peak flowering use registered non-pyrethroid remedies.</p> <p>Weekly spray programme: Preventive treatment: Base application on scouting of lands. Scout 24 plants at random in lands up to 15 ha at weekly intervals from flowering until boll split. Apply FARMAG CYPERMETHRIN when the economic threshold is reached or exceeded i.e. for American (Africa) bollworm an average of 12 eggs or 5 larvae per 24 plants; red bollworm an average of 6 eggs or larvae per 24 plants and spiny bollworm when 2 or more larvae are found during scouting. Stainers and semi-looper will be controlled with applications for control of bollworms.</p> <p>Ground application Application with boom and nozzles. Ensure thorough coverage of the plants. For plants less than 60 cm high apply 100L spray mixture per ha. For taller plants increase the spray mixture volume to 200L per ha. Do not use less than 150 mL FARMAG CYPERMETHRIN per ha on mature cotton.</p> <p>Mistblower Use the lower dosage rate on cotton less than 60cm high and the higher dosage rate on cotton higher than 60cm. Do not use less than 150L FARMAG CYPERMETHRIN on mature cotton.</p> <p>Aerial application Use the lower dosage on cotton less than 60cm high and the higher dosage rate on cotton higher than 60cm. Apply in 30L water per ha.</p> <p>Corrective treatment: Apply as a corrective spray to control an established infestation, i.e. when more than 6 bollworms per 24 plants are found during scouting. The later development stages of the red bollworm already established inside bolls may not be controlled successfully. The success of a treatment can be related to application, density of crop foliage, crop stand and development stages of larvae. Allow up to 4 days for FARMAG CYPERMETHRIN to achieve its full effect. Consider a corrective application as an emergency measure and thereafter the regular programme must be resumed irrespective of the crop stage.</p> <p>Ground application Application with boom and nozzles. Ensure thorough coverage of the plants. For plants less than 60 cm high apply 100L spray mixture per ha. For taller plants increase the spray mixture volume to 200L per ha. Do not use less than 500 mL FARMAG CYPERMETHRIN per ha on mature cotton.</p> <p>Mistblower Use the lower dosage rate on cotton less than 60cm high and the higher dosage rate on cotton higher than 60cm. Do not use less than 500 mL FARMAG CYPERMETHRIN on mature cotton.</p> <p>Aerial application Use the lower dosage on cotton less than 60cm high and the higher dosage rate on cotton higher than 60cm.</p>
GRAPES (Wine and table) Weevils (Snout beetles)	10 mL	Apply as soon as damage occurs. Repeat 3-4 weeks later if necessary. The occurrence of weevils (snout beetles) can be expected from mid-October to mid-November. Refer 'COMPATIBILITY'

CRUCIFERAE American (Africa) bollworm Diamond back moth Thrips	10 mL	Ensure good wetting of the plant by adding a wetting agent. Start spraying at first signs of pest. Repeat every 10-14 days. NOTE: Refer "Warning against American (Africa) bollworm resistance" under 'RESISTANCE WARNING'
GROUNDNUTS American (Africa) bollworm	150 mL/ha	Start spraying as soon as eggs or larvae are noticed. Repeat if necessary. Ground application Apply in 300L water per ha
	150 mL/ha	Aerial application Apply in 30L water per ha Refer "Warning against American (Africa) bollworm resistance" under 'RESISTANCE WARNING'
LUCERNE Lucerne caterpillar	75 mL/ha 75 mL/ha	Apply as full cover spray as soon as pest is noticed. Ground application Apply in 300L water per ha. Aerial application Apply in 30L water per ha.
MACADAMIA NUTS Stink bug	20 mL	Start spraying when nuts are marble size. Repeat after 4 weeks. If necessary, another spray 4 weeks later can be applied, but not later than 90 days before harvest.
MAIZE & SWEETCORN American (Africa) bollworm	1.5mL/100 m plant row	Apply when pest is noticed. Ensure timeous application i.e. not later than 80% beard emergence and before larvae enter beard. Only larvae that come in contact with spray will be controlled. Ground Application If row width allows ground application, apply in not less than 3L water per 100 metre row. Ensure that nozzles are directed towards the maize cobs.
	150 mL/ha	Aerial application Apply in 30L water per ha. Refer "Warning against American (Africa) bollworm resistance" under 'RESISTANCE WARNING'
	3.5 mL/100 m plant row	Preventative treatment against young larvae based on scouting for eggs on plants. Scout fields weekly from 2 to 7 weeks after crop emergence by inspecting at least 100 plants at random per field. Apply 7 days after 2.5% or more plants are found to be infested with eggs. If eggs have hatched, spray at first signs of infestation. Direct spray into the funnel. Repeat application later in season when scouting shows it to be necessary. Larvae that have penetrated the stalk or are longer than 10mm will not be controlled. Ground application Do not apply in less than 3L water per 100 m row. To ensure good coverage inside the funnel as well as on the sides of the plant, use at least two D4/35 or D6/35 nozzles per row.
	350 mL/ha	Aerial application Apply in 30L water per ha.

PEACHES		To determine the volume of water in which FARMAG CYPERMETHRIN should be applied per hectare refer ' GROUND APPLICATION: Volume water to be applied per hectare in deciduous fruit orchards ' above.
American (Africa) Bollworm	5 ml	Apply as full cover spray at 75% petal drop or when pest is noticed. Refer " Warning against American (Africa) bollworm resistance " under ' RESISTANCE WARNING '
Codling moth	5 ml	Apply a full cover spray at 75% petal drop. Repeat at 14-21 day intervals.
False codling moth	5 ml	Early cultivars A full cover spray should be applied 6 weeks before harvest and repeated at 14 day intervals. Late cultivars A full cover spray should be applied 8 weeks before harvest and repeated at 14 day intervals. In the summer rainfall regions first application must not be later than third week December. Canned peaches As above, but last application can be made 3 days before harvest.
Fruit flies	10 ml	All cultivars A full cover spray should be applied 8 weeks before harvest and repeated at 14 day intervals. Canning peaches can be sprayed 3 days before harvest.
Banded fruit weevil (Snout beetle)	10 ml	Apply as soon as damage is noticed. Repeat after 4 weeks if necessary.
PEAS		
American (Africa) Bollworm	150 ml/ha	Ground application Apply in 300-500ℓ water per ha. If necessary repeat.
Lesser army	150 ml/ha	Aerial application Apply in 30ℓ water per ha. Refer " Warning against American(Africa) bollworm resistance " under ' RESISTANCE WARNING '
SORGHUM		
American (Africa) Bollworm	1.5 ml /100m row	Apply when pest is noticed. If bollworms are concealed in the ear, control may be less effective. Later infestation may require a second application.
	150 ml/ha	Ground application Direct application onto ears. Do not apply in less than 3ℓ water per 100 m row.
Stalkborer		Aerial application Apply in 30ℓ water per ha Refer " Warning against American (Africa) bollworm resistance " under ' RESISTANCE WARNING ' Apply at funnel stage when shot hole damage is evident and borer has not yet migrated into the stalks.
	3,5 ml/ha m row	Ground application Do not apply in less than 3ℓ water per 100 m row. To ensure good coverage inside the funnel as well as on the sides of the plant, use at least two D4/35 or D6/35 nozzles per row.
	350 ml/ha	Aerial application Apply in 30ℓ water per ha.
TOMATOES		
American (Africa) Bollworm	15ml	Ground application At first signs of infestation, apply as a full cover spray. Repeat at 7-10 day intervals or as necessitated by crop inspection'
	75 ml/ha	High volume Apply up to 500ℓ per ha spray mixture to plants up to 60cm high and 1 000ℓ per ha or more to plants higher than 60 cm eg. Trellised tomatoes. Do not use more than 150 ml FARMAG CYPERMETHRIN per ha.
	150 ml/ha	Mist Blower Apply in 150 ml water per ha for plants higher than 60 cm. Do not use more than 150 ml FARMAG CYPERMETHRIN per ha. Refer " Warning against American(Africa) bollworm resistance " under ' RESISTANCE WARNING '

PASTURES	150 mL/ha 150mL/ha	Ground application Apply in 300L water per ha. Aerial application Apply in 30L water per ha.
Citrus Thrips (a) Re-infestation 4 to 9 weeks after petal fall following use of any other registered treatment OR (b) Re-infestation mid November to 20 December following use of any other registered treatment.	15 – 20 mL/100 L water 5 – 7 mL/100 L water	<p>Not to be used in orchards where red scale is under biological control.</p> <p>Use progressively lower dosage the later the re-infestation occurs during the designated time period. Apply as a light cover spray to wet foliage canopy to the point of run-off.</p> <p>Use progressively lower dosage the later the re-infestation occurs during the designated time period. Apply as a light cover spray to wet foliage canopy to the point of run-off.</p> <p>NOTE: See use restrictions for citrus at the end of the label.</p>
USE RESTRICTIONS FOR CITRUS:		
<ul style="list-style-type: none"> • Restrict usage to a single application per season and preferable to the period 4 to 9 weeks after petal drop, for thrip control or 5 to 4 weeks before harvest for false codling moth control. Therefore, use FARMAG CYPERMETHRIN only once per season, either for thrip control or false codling moth control. • Restrict usage to orchards in which the red scale (<i>Aonidiella aurantii</i> Mask) is under commercial chemical control. The use of this product may lead to the build-up of red scale. • Where there is any intention to use FARMAG CYPERMETHRIN if thrips reinfestation occurs during the designated time periods, apply a registered preventive treatment for red scale control in the early spring and a second treatment 7 to 11 weeks after petal fall. • Do not use on trees with a history of Tetranychid mite infestations. 		

WAARSKUWINGS

Gevaarstellings: Vlambare vloeistof en damp. Giftig indien ingesluuk. Skadelik in kontak met die vel of indien ingeasem. Veroorsaak velirritasie. Kan respiratoriese irritasie veroorsaak. Mag skade aan organe veroorsaak deur langdurige of herhaalde blootstelling. Baie giftig vir waterlewe. Giftig vir waterlewe met langdurige effekte.

- Onthoudingsperiodes: minimum aantal dae tussen laaste toediening en oes of beweiding:

Appels en pere	14 dae
Bontjies	7 dae
Katoen (beweiding)	28 dae
Koolgewasse	4 dae
Druwe (tafel en wyn)	28 dae
Grondbone (hooi)	7 dae
Lusern (beweiding)	14 dae
Makademia neutra	30 dae
Mielies	14 dae
Weidings	14 dae
Perskes	14 dae
Ertjies	7 dae
Sorghum	28 dae
Suikermielies	14 dae
Tamaties	4 dae
Sitrus.....	28 dae

- **Herbetreding:** Moet nie behandelde gebied betree vir 1 dag na bespuiting nie, tensy beskermende oorklere gedra word.
- In geval van vergiftiging, ontbied 'n geneesheer en toon hierdie etiket aan hom.
- **Lugtoediening:** Stel alle inwoners van die onmiddellike gebied wat bespuiting gaan word, in kennis en reik die nodige waarskuwings uit. Moet nie sput oor of toelaat dat spuitnevel aangrensende gebiede, riviere of damme besoedel nie.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale grond-, klimaats- en bergingstoestande, kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op dié etiket aangedui is nie en die voorkoms van weerstand van die plaag teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effek op mens en dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim deur die verbruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

VOORSORGMAATREËLS

Voorsorgstellings

Algemeen: Indien mediese advies benodig word, hou produkhouer of etiket byderhand. Hou buite bereik van kinders. Lees etiket voor gebruik.

Voorkoming: Hou weg van hitte, warm oppervlaktes, vonke, oop vlamme en ander ontstekingsbronne. Geen rook nie. Hou die houer dig toe. Grond en bind houer en ontvangertoerusting. Gebruik ontplaffingsvaste toerusting. Gebruik gereedskap wat nie vonkel nie. Neem aktie om statiese ontladings te voorkom. Moet nie dampe inasem nie, sput. Was hande, voorarms en gesig deeglik na hantering. Moet nie eet, drink of rook wanneer hierdie produk gebruik word nie. Gebruik slegs buite of in 'n goed geventileerde area. Vermy vrylating na die omgewing. Dra beskermende handskoele/beskermende klere/oogbeskerming/gesigbeskerming.

Reaksie: INDIEN INGESLUK: Ontbied onmiddellik 'n gifsentrum of dokter. INDIEN OP DIE VEL: Was met baie water. Trek onmiddellik alle gekontamineerde klere uit. Spoel vel uit met water/stort. INDIEN INGESEM: Verwyder persoon na vars lug en hou gemaklik om asem te haal. Bel 'n gifsentrum of dokter as jy onwel voel. Kry mediese advies/aandag as jy onwel voel. Spesifieke behandeling (sien aanvullende noodhulpmiddels op hierdie etiket). Spoel mond uit. Indien velirritasie voorkom: Kry mediese advies/aandag. Trek besoedelde klere uit en was dit voor hergebruik. In geval van brand: Gebruik koolstofdioksied (CO₂), droë bluspoeler, skuum om te blus. Versamel gemors.

Berging: Hou houer dig toe. Berg in 'n goed geventileerde plek. Hou koel. Stoer toegesluit.

Wegdoening: Gooi inhoud en houer weg na 'n goedgekeurde afvalverwyderingsaanleg.

- Voorkom die oorwaai van sputnewel na ander gewasse, weiding, riviere, damme en areas nie onder behandeling nie.
- Maak toedieningsapparaat skoon na gebruik. Doe weg met uitspoelwater waar dit nie gewasse, weiveld, riviere, damme en boorgate sal besoedel nie.
- **SPOEL die leë houer DRIEKERUIT soos volg:** Keer die leë houer om oor die sput- of mengtenk en dreineer vir ten minste 30 sekondes nadat die vloeistof drup verlangsaam is. Spoel die leë houer daarna drie keer agtereenvolgens uit met een kwart van die houervolume vars water en gooi die spoelwater in die sput- of mengtenk. Prik die drievoudig uitgespoelde houer deur en gooi weg via 'n goedgekeurde versamaal of herwinner (www.croplife.co.za). Moet nie die houer begrawe, verbrand of skenk aan enige ander partye wat dit as 'n houer vir kos of drank kan gebruik nie.
- Vernietig leë houer deur gate daarin, plat te druk en te begrawe.
- Moet nie leë houer vir enige ander doel gebruik nie.
- Voorkom besoedeling van voedsel, voer, eetgerei en drinkwater.

RELEVANTE STOWWE:

NAAM	PRODUK IDENTIFISEERDER
Supermetriën 97% TS	CAS-Nr.: 52315-07-8
Xileen	CAS-Nr.: 1330-20-7
Tristyrylfenol-etoksilate	CAS-Nr.: 99734-09-5
Kalsiumdodesielbenseensulfonaat	CAS-Nr.: 26264-06-2

SIMPTOME VAN VERGIFTIGING

Veroorsaak branderigheid, jeukerige of tintelende sensasie van die vel wat geredelik binne 24 uur na blootstelling, verdwyn. Inaseming veroorsaak 'n loopneus en krapperie keel. Simptome van innname van 'n hoë dosis sluit naarheid, aanhoudende braking, maagpyn en diarree wat oorgaan in stuutrekings, bewusteloosheid en koma, in.

NOODHULPBEHANDELING

Oë: Spoel oë met skoon, saggloeiende water vir ten minste 20 minute terwyl die boonste en onderste oogledde gereeld opgelig word en tot geen chemikalië teenwoordig is nie. Indien irritasie voorkom en voortduur, ontbied 'n geneesheer.

Vel: Verwyder besoedelde klere en spoel geaffekteerde liggaamsdele deeglik met volop seep en koue water af tot geen teken van chemikalië teenwoordig is nie. Moet nie vel hard vryf nie. Smeer olyfolie aan geaffekteerde dele vir onmiddellike verligting. Kry mediese hulp indien irritasie voorkom of voortduur.

Inaseming: Neem pasiënt na vars lug. Indien asemhalingsprobleme ondervind word of irritasie voorkom en voortduur, kry mediese hulp.

Inname: Moet nie braking veroorsaak nie. Indien die pasiënt by bewussyn is, spoel mond deeglik met water uit. Moet nie iets per mond aan 'n bewusteloze persoon gee nie. Indien pasiënt braak, voorkom inaseming van braaksel. Kry mediese hulp.

NOTA AAN GENEESHEER

Hierdie plaagbeheermiddel bevat 'n piretroid en 'n aromatiese oplosmiddel. Indien plaagbeheermiddel, wanneer ingesluk of tydens braking, in longe ingesasm word, kan ligte tot ernstige chemiese inflammasie van die longe veroorsaak word.

Die maag behoort so spoedig moontlik leeggemaak te word deur die maag versigtig uit te spoel.

Geen teenmiddel is beskikbaar nie. Behandel simptomaties en ondersteunend. In geval van innname, oorweeg maagspoeling met water of 'n 5 % natriumbikarbonaat-oplossing. Behandel stuutrekings met 'n middel daarteen.

WEERSTANDSWAARSKUWING

FARMAG CYPERMETHRIN is 'n groepkode 3 insekdoder. Enige populasie van 'n spesifieke insek mag individue insluit wat 'n natuurlike weerstand teen **FARMAG CYPERMETHRIN** of enige ander groepkode 3 insekdoder het. Indien hierdie insekdoder herhaaldelik aangewend word, kan die weerstandbiedende individue uiteindelik die insek populasie oorheers. Hierdie weerstandbiedende insekte sal waarskynlik nie deur **FARMAG CYPERMETHRIN** of enige ander groepkode 3 insekdoder beheer word nie.

Om weerstand teen insekdoders te vertraag:

- Verminder die eksklusiewe herhaaldelike gebruik van insekdoders met dieselfde groepkode. Wissel af met, of gebruik tenkmengsels van produkte in verskillende insekdodergroep kodes.
- Integreer ander beheermaatreëls (chemies, verbouing, biologies) in insekbeheer programme.

Vir spesifieke inligting oor weerstandsbestuur kontak die registrasiehouer van hierdie produk.

Waarskuwing teen Amerikaanse (Afrika-) bolwurm bestandheid:

Weerstand van die Amerikaanse (Afrika-) bolwurm (*Helicoverpa armigera*) teen sinteriese piretroïede is bevestig.

Katoen: Sintetiese piretroïedes moet slegs gedurende die periode 1 Januarie tot 28 Februarie op katoen toegedien word.

Alle ander gewasse: Moet nie meer as twee toedienings per groeiseisoen toedien nie.

As 'n piretroïedbespuiting swak beheer gee moet nie met enige sintetiese piretroïed, selfs teen 'n korreklike dosis, herspuit nie. Gebruik 'nprodukt uit 'n ander chemiegroep.

WYSE VAN AKSIE:

Werk op die senuwestelsel van die insek, en versteur die funksie van neurone deur interaksie met die natriumkanaal. Nie-sistemiese insekdoder met kontak- en maagwerkung. Toon ook anti-voeding aksie. Goeie nawerking op behandelde plante.

GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui.

VERENIGBAARHEID

- Moet nie **FARMAG CYPERMETHRIN** met seewierekstrak, varmidotin, protofos, chinometionaat of amitras bevattende plaagbeheermiddels meng nie.
- Moet nie 'n benattingsmiddel by die **FARMAG CYPERMETHRIN** spuitmengsel voeg wanneer appels, pere en druive gespuit word nie.
- **FARMAG CYPERMETHRIN** is vereenbaar met 'n wye verskeidenheid plaagbeheermiddels. Dit is nie vir Cipla Agricare (Pty) Ltd moontlik om alle kombinasies te toets nie en word dit aanbeveel om 'n vereenigbaarheids toets op klein skaal uit te voer indien daar enige twyfel bestaan.
- Die doeltreffendheid van **FARMAG CYPERMETHRIN** kan negatief deur 'n hoog pH waarde van die spuitmengsel beïnvloed word. Sipermetriën, die aktiewe bestanddeel in **FARMAG CYPERMETHRIN**, is meer stabiel by 'n pH van ongeveer 4. Indien 'n versuringsmiddel gebruik word om die pH te verlaag, moet dit eers met die verduningswater gemeng word, alvorens **FARMAG CYPERMETHRIN** bygevoeg word.
- Byvoeging van melasse by die spuitmengsel versuur en verminder die verdamping van die spuitnewel. Vir grontoediening behoort 10 % melasse per volume bygevoeg te word en vir lugbespuiting 20 %.

MENGINSTRUKSIES

Vul spuitenk half met skoon water. Voeg die verlangde hoeveelheid **FARMAG CYPERMETHRIN** by terwyl deeglik geroer word. Vul die tenk hierna met water tot die verlangde volume en hou vol om mengsel in tenk te roer totdat bespuiting voltooi is. Lees inligting onder 'VERENIGBAARHEID' hierbo aangaande pH van spuitmengsel.

GRONDOEDIENING

Wanneer 'n konvensionele hoog volume spuitapparaat en newelblaser gebruik word, gebruik holkeëlspuipunte wat 'n medium tot fyn druppelgrootte lewer. Die spuitapparaat moet akkuraat gekalibreer word en tydens toediening moet daar seker gemaak word van goeie bedekking van gewas en eweredige verspreiding van spuitmengsel.

Volume water wat per hektaar in sagte vrugteboorde toegedien moet word:

Die volume water wat per hektaar in sagte vrugteboorde toegedien moet word, word volgens die Boomryvolume-formule soos deur Unrath beskryf, bereken:

Die formule is as volg:

$$\text{Boom hoogte} \times \text{boom diameter} \times 937 = \text{Volume water/ha wat toegedien moet word}$$

Ry wydte

Die afmetings wat in die formule gebruik word, is dié van die boom met volle blaarbedekking. Aangesien sagtewrugtebome nie altyd ten volle met blare bedek is nie, is die volume water wat per hektaar toegedien moet word tydens die vroeë groestadia van die bome, as volg:

Van groenpunt tot 30 % blom - 60 % van die HV wat toegedien moet word

Van ca. volblom tot ca. November - 80 % van die HV wat toegedien moet word

Van ca. begin Desember - 100 % van die HV wat toegedien moet word

Wanneer plaagbeheermiddels as lae volume bespuitings toegedien word, moet die korrekte hoeveelheid plaagbeheermiddel, soos bereken vir hoog volume bespuitings, per hektaar toegedien word.

LUGTOEDIENING

FARMAG CYPERMETHRIN kan slegs deur 'n geregistreerde Lugbespuitingsoperator met 'n korrek gekalibreerde, geregistreerde vliegtuig volgens die instruksies van SANS 10118: 2009 (The Aerial Application of Pesticides) uit die lug bespuit word. Verseker dat die sputtmengsel eweredig oor die teikenarea versprei word, en die verlies aan sputtmengsel tydens toediening tot 'n minimum beperk word. Dit is daarom belangrik om aan die volgende vereistes te voldoen:

- **Volume:** 'n Sputtmengsel volume van 30 liter per hektaar word aanbeveel. Hierdie produk is nie teen 'n verlaagde volume getoets nie. Die registrasiehouer kan nie effektiwiteit waarborg, of verantwoordelik gehou word vir enige nadelige effekte indien hierdie produk teen 'n laer volume, as hierbo aanbeveel, toegedien word nie.
- **Druppel bedekking:** 30 tot 40 druppels per cm² moet op die teikenarea herwin word.
- **Druppelgrootte:** 'n Druppelspektrum met 'n VMD van 250 tot 280 mikrons word aanbeveel. Beperk die produksie van druppels kleiner as 150 mikrons (hoë drywing en verdampingspotensiaal) tot 'n minimum.
- **Vlieghoogte:** Handhaaf die hoogte van die sputtbalk bo die teiken op 3 tot 4 meter. Moet nie sputt wanneer die vliegtuig duik nie, uitklim of draai nie.
- Gebruik geskikte atomiseringsapparaat wat die vereiste druppelgrootte en bedekking sal produseer, maar die minste verlies van produk verseker. Die sputstelsel moet 'n druppelspektrum met die kleinste moontlike Relatiewe Span produseer.
- Plaas al die atomiseerders in die binnekste 60 tot 75 % van die vlerkspan om te verhoed dat druppels binne-in die vlerkpuntvorteeks beweeg.
- Die verskil in temperatuur tussen die nat- en droëboltermometer van 'n swaaihigrrometer, moet nie 8 °C oorskry nie.
- Stop bespuiting indien die windspoed 15 km/uur oorskry.
- Stop bespuiting tydens turbulente, onstabiele en droë toestande gedurende die hitte van die dag.
- Bespuiting onder temperatuur inversie toestande (deur bo of binne die inversie laag te sputt) en/of hoë lugvog toestande (relatiewe humiditeit 80 % en meer) mag tot volgende probleme aanleiding gee:
 - verlaagde effektiwiteit aangesien die druppels as 'n wolk in die lug bly hang en moontlik verdamp (onvoldoende bedekking op teiken).
 - skade aan nie-teiken gewasse of sensitiewe areas as gevolg van wegdrywing van die sputtwolk na nie-teiken area.
- Verseker dat die Lugbespuitingsoperator presies weet watter lande bespuit moet word.
- Dit is noodsaaklik om 'n versekering van die Lugbespuitingsoperator te verkry dat aan al die bogenoemde vereistes voldoen sal word en dat data van belang in 'n logboek saamgevat is vir toekomstige verwysing.

GEWAS/PLAAG	DOSIS / 100ℓ WATER OF SOOS AANGEDUI	OPMERKINGS
BOSBOU		
Denne pou-oogmot larf	50 mL/ha	Lugtoedienving Dien in 30 ℓ water per ha toe.
Wilger pou-oogmot larf	100 mL/ha	Lugtoedienving Dien in 30 ℓ water per ha toe.
Wattleboom sakwurm	100 mL/ha	Lugtoedienving Dien in 30 ℓ water per ha toe. Begin behandeling in November/Desember.
ALLEGEWASSE		
Snywurms		Grondtoedienving Algehele bespuiting: dien in 300ℓ water per ha toe. Bybespuiting (300 mm breed): dien in ten minste 3ℓ water per 100 m ry toe.
	100 mL/ha 0.33 mL/100m plantry	
	100 mL/ha	Lugtoedienving Dien in 30ℓ water per has toe

APPELS & PERE		
Koddingmot Blaarroller	5 mL	Om die volume water waarin FARMAG CYPERMETHRIN per has toegedien moet word, te bepaal, lees " GRONDOEDIENING: Volume water wat per hektaar in sagtevrugteboorde toegedien moet word ". Lees ook " VERENIGBAARHEID ".
Amerikaanse (Arika-) Bolwurm	5 mL	Dien eerste bespuiting met 75% blomblaarval toe en herhaal elke 14 dae. Dien as 'n korrektiwe behandeling toe maar nie vroeër as 75% blomblaarval nie. Lees " Waarskuwing teen Amerikaanse (Afrika) bolwurm bestandheid " onder " WEERSTANDWAARSKUWING "
Gebande vrugte-kalander (Snuitkewer)	10 mL	Die twee bespuitings toe, die eerste tydens 75% blomblaarval en herhaal 4 weke later. 'n Verdere hoe of lae volume bespuiting kan middel Januarie toegedien word om laatseisoen skade te voorkom.
BOONTJIES Amerikaanse (Arika-) bolwurm	150 mL/ha	Grondtoediening Dien in 300 L water per ha toe. Verken landerye en indien nodig herhaal bespuiting.
	150 mL/ha	Lugtoediening Dien in 30 L water per ha toe. Lees " Waarskuwing teen Amerikaanse (Afrika-) bolwurm bestandheid " onder " WEERSTANDWAARSKUWING "

KATOEN Amerikaanse (Afrika) rooi-en-stekelbolwurms Vlekbesies Vals-(Plusia) landmeter		<p>Lees "Waarskuwing teen Amerikaanse (Afrika-) bolwurm bestandheid" onder "WEERSTANDWAARSKUWING". Om te voldoen aan die beginsels van plaagbestuur en geïntegreerde beheermetodes word aanbeveel om FARMAG CYPERMETHRIN gedurende die vrugvormingperiode, vanaf hoofblom tot bolbars te gebruik, d.w.s. vanaf ongeveer 10 tot 22 weke na opkom van die gewas. Die gebruik van FARMAG CYPERMETHRIN is hoofsaklike bedoel as 'n voorkomedende Eheermaatreël teen al die bolwurms gebaseer op gereelde weeklikse verkenning van landerye of 'n gereelde sputiprogram met 7-dae tussenposes. Gewoonlik neem 'n sputiprogram aanvang aan die begin van die blomstadium, d.w.s. omtrent 6 weke na opkom an die gewas. Van dan tot hoofblomperiode moet geregistreerde nie piretroïed middels gebruik word.</p> <p>Weeklikse sputiprogram: Voorkomende behandeling: Baseer toediening op verkenning van landerye. Verken 24 FARMAG CYPERMETHRIN toe wanneer ekonomiese drempel bereik of oorskry word en is vir Amerikaanse (Afrika-) bolwurm, 'gemiddeld van 12 eiern of 5 larwes per plante; vir rooibolwurm, 'n gemiddeld van 6 eiern of 2 larwes op 24 plante gedurende verkenning gevind word. Vlekbesies en valsslandmeter sal ook beheer word met bolwurmsbespuittings.</p> <p>75 mL</p> <p>Grondtoediening Toediening met sputtbalk en -punte. Dra sorg dat die plante deeglik bedek word. Gebruik 100ℓ sputmengsel per ha op lante korter as 60cm. Vir groter plante vermeerder die hoeveelheid sputmengsel geleidelik tot 200ℓ sputmengsel per ha. Moet nie minder as 150 mL FARMAG CYPERMETHRIN per ha op volwasse katoen gebruik nie.</p> <p>Newelblaser Gebruik die laer dosis op katoen korter as 60cm en die hoër as 60 cm. Moet ni minder as 150 mL FARMAG CYPERMETHRIN per ha op volwasse katoen toedien nie.</p> <p>87 mL of 175 mL/ha</p> <p>Lugtoediening Dien in 30ℓ water per ha toe. Gebruik die laer dosis op katoen korter as 60cm en die hoër dosis op plante hoer as 60cm.</p> <p>Korrektiewe behandeling: Dien as korrektiewe bespuiting toe om 'n bestaande infestasie te beheer, d.w.s. waar meer as 6 bolwurms per 24 plante tydens verkenning gevind word. Die latere ontwikkelingsstadia van die rooibolwurm wat reeds in bolle gevlest is, sal nie effekief beheer word nie. Die sukses van 'n behandeling hang af van die toediening, digtheid van gewaslaaf, stand van plante en van die ontwikkelingsstadia van die larwes. Laat 4 dae toe vir FARMAG CYPERMETHRIN om sy volle effek te bereik. 'n Korrektiewe toediening moet as 'n noodmaatreël beskou word en daarna moet met 'n gereelde program voortgegaan word ongeag die ontwikkelingsstadium van die katoen.</p> <p>250 mL</p> <p>Grondtoediening Toediening met sputtbalk en punte. Dra sorg dat die plante deeglik bedek word. Gebruik 100ℓ sputmengsel per ha op plante korter as 60cm. Vir groter plante vermeerder die hoeveelheid sputmengsel geleidelik tot 200ℓ sputmengsel per ha. Moet nie minder as 500 mL FARMAG CYPERMETHRIN per ha op volwasse katoen toedien nie.</p> <p>Newelblaser Gebruik die laer dosis op plante korter as 60 cm en die hoër dosis op lante hoër as 60 cm. Moet nie minder as 500 mL FARMAG CYPERMETHRIN op volwasse katoen toedien nie.</p> <p>250 or 500 mL/ha</p> <p>Lugtoediening Gebruik die laer dosis op plante korter as 60cm en die hoër dosis op lante hoër as 60cm. Moet nie minder as 500 mL FARMAG CYPERMETHRIN op volwasse katoen toedien nie. Die toe in 30ℓ water per ha.</p> <p>DRUIWE (wyn en tafel) Kalanders (snuitkewers)</p> <p>10 mL</p> <p>Dien sodra skade opgemerk word toe. Herhaal 3-4 weke later indien nodig. Kalanders (Snuitkewers) kom gewoonlik tussen middel Oktober tot middel November voor. Lees "VERENIGBAARHEID"</p>
---	--	--

KOOLGEWASSE Amerikaanse (Afrika-) bolwurm Ruitrugmottarf Blaaspootjies	10 ml	Verseker goeie benetting van plante. Begin spuit met eerste tekens van plaag. Herhaal elke 10-14 dae. Lees "Waarskuwing teen Amerikaanse (Afrika-) bolwurm bestandheid" onder "WEERSTANDWAARSKUWING"
GRONDBONE Amerikaans (Afrika-) bolwurm	150 ml/ha 150 ml/ha	Begin spuit sodra eiers of larwes opgemerk word. Herhaal indien nodig. Grondtoediening Dien in 300 ℥ water per ha toe. Lugtoediening: Dien toe in 30ℓ water per ha. Lees "Waarskuwing teen Amerikaanse (Afrika-) bolwurm bestandheid" onder "WEERSTANDWAARSKUWING"
LUSERN Lusernruspe	75 ml/ha 75 ml/ha	Dien as 'n voldekkende bespuiting toe sodra plaag opgemerk word. Grondtoediening Die toe in 300ℓ water per ha Lugtoediening Die toe in 300ℓ water per ha
MAKADAMIANEUTE Stinkbesie	20 ml	Begin spuit wanneer neutre albastergrootte bereik en herhaal na 4 weke. Indien nodig, kan 'n verder bespuiting 4 weke later toegedien word. Maak egte seker dat daar ten minste 90 dae tussen hierdie laaste bespuiting en oes is.
MIELIES & SUIKERMIELIES Amerikaanse (Afrika-) bolwurm	1.5 ml / 100 m ry 150 ml / ha 3,5 ml / 100 m ry 350 ml / ha	Dien toe wanneer plaag opgemerk word. Sorg dat bespuiting betyds gedoen word, dit is, nie later as 80% baardverskyning nie en voordat die larwes die baard binnedring. Slegs larwes wat in aanraking met die sproei kom, sal beheer word. Grondtoediening Indien rywydte grondtoediening toelaat, gebruik in ten minste 3ℓ water per 100 m ry. Dra sorg dat die spuitpunte na die mieliekoppe gering is. Lugtoediening Dien toe in 30ℓ water per ha. Lees "Waarskuwing teen Amerikaanse (Afrika-) bolwurm bestandheid" onder "WEERSTANDWAARSKUWING" Voorkomende behandeling teen jong larwes gebaseer op verkenning vir eiers op plante. Verken landerye elke week van 2 tot 7 weke na opkom van gewas deur ten minste 100 plante ewekansig per landery te ondersoek. Spuit 7 dae nadat eiers op 2,5% of meer van die plante gevind is. Indien die eiers al uitgebroei het, spuit sodra eerste tekens van infestasie opgemerk word. Rig bespuiting direk in die kelk in. Herhaal die toediening later in die seisoen indien verkenning aandui dat dit nodig is. Larwes wat reeds die stam binnegedring het of langer as 10mm is, sal nie beheer word nie. Grondtoediening Moet nie in minder as 3ℓ water per 100m ry toedien nie. Om seker te maak dat goeie bedekking binne in die kelk in sowel as aan die sye van die plant verky word, gebruik ten minste twee D4/35 of D6/35 spuitpunte oor 'n ry. Lugtoediening Dien toe in 30ℓ water per ha.

PERSKES		Om die volume water waarin FARMAG CYPERMETHRIN per ha toegedien moet word, te bepaal, lees "GRONDOEDIENING: Volume water wat per hektaar in saagte vrugteboorde toegedien moet word".
Amerikaanse (Afrika-) bolwurm	5 ml	Dien toe as 'n voldekkbespuiting by 75% blomblaarval van wanneer plaag opgemerk word. Lees "Waarskuwing teen Amerikaanse (Afrika-) bolwurm bestandheid" onder "WEERSTANDWAARSKUWING"
Kodlingmot	5 ml	Dien 'n volbedekkende bespuiting toe by 75% blomblaarval. Herhaal met 14-21 dae tussenposes.
Valskodlingmot	5 ml	Vroeëkultivars Dien toe as 'n voldekkbespuiting 6 weke voor oes en herhaal met 14 dae tussenposes. Laat kultivars Dien toe as 'n voldekkbespuiting 8 weke voor oes en herhaal met 14 dae tussenposes In die somerreënvalgebied moet die eerste behandeling nie later as die derde week in Desember toegedien word nie. Inmaakperskes Soos hierbo, behalwe dat laaste 3 dae voor oes gespuit kan word. Alle Kultivars Dien 'n voldekkbespuiting 8 weke voor oes toe en herhaal met 14 dae tussenposes. Ingelede perskes kan tot 3 dae voor oes gespuit word.
Vrugtevlieë	10 ml	
Gebande vrugte-kalander (Snuikewer)	10 ml	Dien toe sodra skade opgemerk word en herhal indien nodig na 4 weke
ERTJIES		Grondtoediening
Amerikaans (Afrika-) bolwurm	150 mℓ/ha	Dien toe in 300 – 500 ℓ water. Herhaal indien nodig
Klein kommandowurm	150 mℓ/ha	Lugtoediening: Dien toe in 30ℓ water per ha. Lees "Waarskuwing teen Amerikaanse (Afrika-) bolwurm bestandheid" onder "WEERSTANDWAARSKUWING"
SORGHUM		Spuik sodra plaag opgemerk word. Indien bolwurms binne versteek is, mag die beheer minder effekief wees. Latere infestasies mag 'n tweede bespuiting vereis.
Amerikaanse (Afrika-) bolwurm	1,5 mℓ / 100 m ry	Grondtoediening Rig toediening op are. Moet ni minder as in 3ℓ water per 100 m ry toedien nie.
Stronkboorder	150 mℓ / ha	Lugtoediening Dien toe in 30ℓ
	3,5 mℓ / ha m ry	Lees "Waarskuwing teen Amerikaanse (Afrika-) bolwurm bestandheid" onder "WEERSTANDWAARSKUWING" Dien toe by kelkstadium wanneer haelskade waargeneem word en boorder nog nie in die stam inbeweeg het nie. Grondtoediening
	350 mℓ/ha	Moet nie in minder as 3ℓ water per 100 m ry toedien nie. Om seker te maak dat goeie bedekking binne in die kelk en sowel as aan die sye van die plant verkry word, gebruik ten minste twee D4/35 of D6/35 spuitpunte oor 'n ry. Lugtoediening Dien toe in 30ℓ water per ha.

TAMATIES Amerikaanse (Afrika-) bolwurm	15 mL	Grondtoediening Dien as 'n voldekkbespuiting toe met eerste teken van 'n infestasie. Herhaal met 7-10 dae tussenposes of soos nodig geag word met gewasinspeksie.
	75 mL/ha	Hoë volume Dien tot 500ℓ sputtmengsel toe op plantte korter as 60cm en in 1 000ℓ of meer sputtmengsel toe op plantte hoër as 60 cm by opgeleide tamaties. Moet nie meer as 150 mL FARMAG CYPERMETHRIN per has toedien nie.
	150 mL/ha	Newelblaser Dien toe in 150ℓ water per ha op plantte 60cm en korter. Dien toe in 500ℓ water per ha op plantte 60cm en hoër. Moet nie meer as 150mℓ FARMAG CYPERMETHRIN per ha toedien nie. Lees "Waarskuwing teen Amerikaanse (Afrika-) bolwurm bestandheid" onder "WEERSTANDWAARSKUWING"
WEIDING Kommandowurm	150 mL/ha	Grondtoediening Die toe in 300ℓ water per ha
	150 mL/ha	Lugtoediening Dien toe in 30ℓ water per ha
SITRUS Blaaspootjies (a) Herinfestasie 4 tot 9 weke na blomblaarval na gebruik van enige ander geregistreerde behandeling OF (b) Herinfestasie middel November tot 20 Desember na gebruik van enige ander geregistreerde behandeling.	15 – 20 mL/100 ℓ water	Nie vir gebruik in boorde waar rooidopluis onder biologiese beheer is nie. Gebruik progressief laer dosisse hoe later herinfestasie voorkom gedurende die aangeduide periode. Dien as 'n ligte dekbespuiting toe. Benat blaardak tot die punt van afloop.
	5 – 7 mL/100 ℓ water	Gebruik progressief laer dosisse hoe later herinfestasie voorkom gedurende die aangeduide periode. Dien as 'n ligte dekbespuiting toe. Benat blaardak tot die punt van afloop. Nota: Sien gebruiksbeperkinge by sitrus onderaan tabel.
GEBRUIKSBEPERKINGE BY SITRUS		
<ul style="list-style-type: none"> • Beperk toediening tot 'n enkelle bespuiting per seisoen en verkiesslik in die periode 4 tot 9 weke na blomblaarval vir blaaspootjiebeheer of 5 tot 4 weke voor oes vir vlagskoddingmot beheer. Dien dus FARMAG CYPERMETHRIN slegs eenkeer per seisoen toe vir of blaaspootjieof vlagskoddingmot beheer. • Beperk gebruik tot boorde waar rooidopluis (<i>Anonidella auranti Mask</i>) onder kommersiele chemiese beheer is. Die gebruik van hierdie produk vir blaaspootjiebeheer mag die opbou van rooidopluis tot gevolg he. • Waar daar enige voorname is om FARMAG CYPERMETHRIN te gebruik indien blaaspootjie besmetting voorkom gedurende die aangeduide periodes, dien 'n geregistreerde voorkomende behandeling vir rooidopluisbeheer vroeg in die lente en 'n tweede behandeling 7 tot 11 weke na blomblaarval toe. Moet nie op bome gebruik met 'n geskiedenis van teranychid mynt infestasie nie. 		

FARMAG

CYPERMETHRIN

Reg. No. / Nr L3207 Act No. 36 of 1947/ Wet Nr 36 van 1947



INSECTICIDE

**READ THE LABEL AND ATTACHED PACKAGED LEAFLET BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

IRAC INSECTICIDE GROUP CODE

3

IRAC INSEKDODER GROEPKODE

An emulsifiable concentrate contact and stomach insecticide for the control of agricultural pests as indicated.

'n Emulgeerbare konsentraat kontak- en maaginsekodder vir die beheer an landbouplae soos aangedui.

ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL:

Cypermethrin (pyrethroid)200 g/l..... Sipermetrien (piretroïed)



DANGER

Batch No:

D.O.M:

D.O.E:

Net Mass:

See reverse of packaging for details.

FOR FULL PARTICULARS SEE ATTACHED PAMPHLET.
VIR VOLLEDIGE BESONDERHEDE KYK AANGEHEGTE ETIKET.

UN No. / VR Nr: 3351



REGISTRATION HOLDER/ REGISTRASIEHOUER
Castle Ag-Chem (Pty) Ltd.
Reg. No. / Nr. 2007/033120/07
61 Marshall Drive, Old Mill Industrial Park,
Mount Edgecombe.
Tel: (031) 003 3486

Hazard Statements:

Flammable liquid and vapour.

Toxic if swallowed.

Harmful in contact with skin or if inhaled.

Causes skin irritation.

May cause respiratory irritation.

May cause damage to organs through prolonged or repeated exposure.

Very toxic to aquatic life.

Toxic to aquatic life with long lasting effects.

Precautionary Statements:

Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking.

Do not breathe vapours, spray.

IF SWALLOWED: Immediately call a poison center or doctor.

IF ON SKIN: Wash with plenty of water.

IF INHALED: Remove person to fresh air and keep comfortable for breathing.

If skin irritation occurs: Get medical advice/attention.

In case of fire: Use carbon dioxide (CO₂), dry extinguishing powder, foam to extinguish.

Poison Information Helpline (24 hrs): 0861 555 777



BARCODE

100450

