

## WARNINGS

**Hazard Statements:** May be harmful if swallowed. May be harmful in contact with skin. Causes mild skin irritation. May cause an allergic skin reaction. Causes serious eye damage. Toxic if inhaled. May cause damage to organs through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects.

### Pre-Harvest Interval:

Dry beans, Green beans and Soybeans..... 14 days  
Groundnuts.....56 days

**NOTE:** Withholding periods above are based on local requirements for residue levels and may differ from other countries to which treated products may be exported. If the treated crop is intended for export, please consult the relevant exporting body regarding the pre-harvest interval requirements of the countries to which the produce will be exported.

- **FARMESAFEN 250 SL** has been tested on the most important commercial varieties without any phytotoxic effects. More susceptible cultivars may be released in future which will need to be tested; for this, apply **FARMESAFEN 250 SL** to a limited test area before commercial applications are made.
- Store in the original container in a safe place away from food and feed.
- Keep out of reach of children, animals and uninformed persons.
- **Re-entry Interval**– Do not enter the treated area within 1 day after application unless wearing suitable protective clothing.

**Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be effective under all conditions. The activity and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the weed against the remedy, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal, or for lack of performance of the remedy concerned due to failure by the user to follow the label instructions, or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.**

## PRECAUTIONS:

### Precautionary Statements

**General:** If medical advice is needed, have product container or label at hand. Keep out of reach of children. Read label before use.

**Prevention:** Use only outdoors or in a well-ventilated area. Contaminated work clothing should not be allowed out of the workplace. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection.

**Response:** IF ON SKIN: Wash with plenty of soap and water. IF INHALED: Remove person to fresh air and keep comfortable for breathing. Immediately call a doctor. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Get medical advice/attention if you feel unwell. If skin irritation occurs: Get medical advice/attention. If skin irritation or rash occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse. Collect spillage.

**Storage:** Store in a well-ventilated place. Keep container tightly closed. Store locked up.

**Disposal:** Dispose of contents and container to an approved waste disposal plant.

- Do not smoke, eat or drink whilst handling this product.
- Prevent contamination of food, feed, drinking water as well as eating utensils.
- **TRIPLE RINSE the empty container as follows:** Invert the empty container over the spray or mixing tank and drain for at least 30 seconds after the flow has slowed down to dripping. Thereafter rinse the empty container three times in succession with one quarter of the container volume fresh water and decant the rinsate into the spray or mixing tank. Puncture the triple rinsed container and dispose of via an approved collector or recycler ([www.croplife.co.za](http://www.croplife.co.za)). Do not bury, burn or donate the container to any other parties that may use it as a container for food or beverages.
- Destroy empty container and do not use for any other purpose.
- Whilst spraying, avoid contact with the spray as much as possible. Avoid spray drift onto other crops, grazing, rivers or dams.
- Clean the applicator thoroughly following use and dispose of the wash water where it will not contaminate crops, grazing, rivers and dams.
- Destroy the empty container and do not use it for any other purpose.

# FARMESAFEN 250 SL

Reg. No. / Nr L10319 Act No. 36 of 1947/ Wet Nr 36 van 1947

HERBICIDE



**READ THE LABEL AND ATTACHED PACKAGED LEAFLET BEFORE USE  
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

HRAC HERBICIDE GROUP CODE

14

ONKRUIDDODER GROEPKODE

A post-emergence water soluble concentrate selective herbicide for the selective control of various annual broadleaf weeds in the crops as listed.

'n Na-opkoms water oplosbare konsentraat selektiewe onkruiddoder vir die selektiewe beheer van verskeie eenjarige breëblaaronkruid in gewasse soos aangedui.

## ACTIVE INGREDIENT/AKTIEWE BESTANDEDEEL:

Fomesafen..... 250 g/ ℓ



**DANGER**

### Hazard Statements:

May be harmful if swallowed.  
May be harmful in contact with skin.  
Causes mild skin irritation.  
May cause an allergic skin reaction.  
Causes serious eye damage.  
Toxic if inhaled.  
May cause damage to organs through prolonged or repeated exposure.  
Very toxic to aquatic life with long lasting effects.

### Precautionary Statements:

Do not breathe spray, vapours.  
IF ON SKIN: Wash with plenty of soap and water.  
IF INHALED: Remove person to fresh air and keep comfortable for breathing. Immediately call a doctor.  
IF IN EYES: Rinse cautiously with water for several minutes. Immediately call a doctor.  
If skin irritation or rash occurs: Get medical advice/attention.

Batch No:

D.O.M:

D.O.E:

Net Mass:

See reverse of packaging for details.

FOR FULL PARTICULARS SEE ATTACHED PAMPHLET.  
VIR VOLLEDIGE BESONDERHEDE KYK AANGEHEGTE ETIKET.

UN No. / VR Nr: 3082

Poison Information Helpline (24 hrs): 0861 555 777



REGISTRATION HOLDER/ REGISTRASIEHOUER  
Cedar Falls Properties 130 (Pty) Ltd  
Reg. Nr.: Reg. No. 2006/031383/07  
61 Marshall Drive, Old Mill Industrial Park,  
Mount Edgecombe, South Africa, 4300  
Tel: 031 003 3486



BARCODE

## RELEVANT SUBSTANCES

NAME	PRODUCT IDENTIFIER
Fomesafen TC	CAS-No.: 72178-02-0
Alkyl polyglucoside	CAS-No.: 68515-73-1
Calcium dodecylbenzenesulfonate	CAS-No.: 26264-06-2
1,2-Benzisothiazolin-3-one	CAS-No.: 2634-33-5

## SYMPTOMS OF HUMAN POISONING

Headache, dizziness, nausea.

## FIRST AID TREATMENT

Do not induce vomiting. Seek medical advice.

## NOTE TO PHYSICIAN

Symptomatic treatment.

## RESISTANCE WARNING

For resistance management, **FARMESAFEN 250 SL** is a group code 14 herbicide. Any weed population may contain individuals naturally resistant to **FARMESAFEN 250 SL** and other group code 14 herbicides. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds may not be controlled by **FARMESAFEN 250 SL** or any other group code 14 herbicide.

- To delay herbicide resistance:
  - Avoid exclusive repeated use of herbicides from the same herbicide group code.
  - Alternate or tank mix with registered products from different herbicide group codes.
  - Integrate other control methods (chemical, cultural, biological) into weed control programmes.
- For specific information on resistance management contact the registration holder of this product.
- The presence of resistant weeds is difficult to detect prior to use and therefore neither the Registration Holder nor the Distributor can accept responsibility for any losses as a result of **FARMESAFEN 250 SL** not controlling resistant weeds. Do not apply **FARMESAFEN 250 SL** alone in situations where weed resistance has been confirmed. Sub-standard application of **FARMESAFEN 250 SL** will lead to inferior weed control and may assist in development of herbicide resistance.

## MODE OF ACTION:

Protoporphyrinogen oxidase inhibitor. Selective herbicide, absorbed by both leaves and roots, with very limited translocation in the phloem.

## USE RESTRICTIONS:

1. **FARMESAFEN 250 SL** should not be used where soybeans, dry beans, green beans or groundnuts are inter-planted with other crops.
2. All other crops, except for those on which the product is registered, should not be planted within 12 months of treatment.
3. Do not apply **FARMESAFEN 250 SL** to any crop or situation not mentioned in this label.
4. **FARMESAFEN 250 SL** must not be used in the North West Province or the Western Free State.
5. **FARMESAFEN 250 SL** can only be used on the following cultivars:
  - Soybeans - All already existing cultivars.
  - Dry beans - Nuweveld, Teebus and Bonus, World Cup, OPS RS4
  - Green beans - Strike, Provider, Cape, Derby, Palati, Newton
  - Groundnuts - Sellie.
6. Do not apply this product in combination or tank mixture with any other product or agricultural remedy which is not specifically listed on this label. Cedar Falls Properties 130 (Pty) Ltd accepts no responsibility for any negative effects experienced if the instructions on this label are not adhered to, unless a Meridian representative was consulted beforehand.

## **DIRECTIONS FOR USE: Use only as directed**

**FARMESAFEN 250 SL** is a selective post-emergence broadleaf herbicide that can be applied up to the pre-bloom stage of the crop but must not be applied later than 4 weeks after planting and not later than the 4 leaf stage of the weeds.

Application at a later growth stage may increase the chance of some re-growth from treated weeds. It is advisable to wait until the majority of broadleaf weeds have germinated before spraying with **FARMESAFEN 250 SL** (see weed list below).  
Rainfall after 4 - 6 hours after application should not affect the activity of **FARMESAFEN 250 SL**.

If the normal cultivation method is the use of pre-emergence herbicides, then this practice should be continued followed by **FARMESAFEN 250 SL** for later weed germination.

Temporary crop damage symptoms, circular scorch marks on leaves and possible retardation of growth may occur after a **FARMESAFEN 250 SL** treatment but will not result in yield losses.

### **COMPATIBILITY**

**FARMESAFEN 250 SL** should not be mixed with other products.

**FARMESAFEN 250 SL** may be used in sequence with other registered herbicides provided that a sufficient time interval is allowed so that the other weeds (such as grasses) are not stressed. For example, where **FARMESAFEN 250 SL** is used before or after an application of fluzafop-p-butyl containing products, allow 14 days between spraying.

When using other products in a spray program with **FARMESAFEN 250 SL**, always consult the labels of those particular products regarding directions for their application and crop rotation recommendations.

### **CLIMATIC CONDITIONS:**

**FARMESAFEN 250 SL** should be applied when weeds are actively growing, under favourable temperature and moisture conditions. **FARMESAFEN 250 SL** should not be applied when crops and weeds are under drought stress or cold temperatures, when poor weed control will result.

### **SOIL BORNE DISEASES:**

Do not apply **FARMESAFEN 250 SL** to soybeans, green beans, groundnuts or dry beans which already show symptoms of soil borne diseases. This is because the combination of a **FARMESAFEN 250 SL** treatment and disease stress may result in stand losses.

### **APPLICATION METHODS**

Ensure that the equipment is correctly calibrated and is checked regularly during application to ensure even and accurate application. Do not apply **FARMESAFEN 250 SL** if rain is expected within 4-6 hours. **FARMESAFEN 250 SL** can be applied by ground equipment. Use only clean water.

### **GROUND APPLICATION**

Apply **FARMESAFEN 250 SL** in a minimum of 200 - 300 l water/ha. Use flat fan or hollow cone nozzles and a minimum spray pressure of 150kPa. Ensure that spray mixture is evenly distributed to give good coverage of the weeds.

### **RECOMMENDATIONS**

<b>CROP</b>	<b>FARMESAFEN 250 SL ℓ /ha</b>	<b>REMARKS</b>
<b>Soybeans, Dry beans, Green Beans, Groundnuts.</b> Broadleaf weeds only	1 ℓ / ha	Spray when the weeds are between the seedling and 4-leaf stages and pre-blossom of the crop. See below for the weeds controlled

**WEEDS NORMALLY CONTROLLED:**

The following broadleaf weeds (seedling up to the 4 leaf stage) will be controlled by FARMESAFEN 250 SL at 1 ℓ / ha:

<i>Amaranthus hybridus</i>	Cape pigweed
<i>Datura ferox</i>	large thorn apple
<i>Galinsoga parviflora</i>	gallant soldier
<i>Portulaca oleracea</i>	purslane
<i>Schkuhria pinnata</i>	dwarf marigold
<i>Sida cordifolia</i>	heartleaf sida
<i>Richardia brasiliensis</i>	Tropical Richardia
<i>Acanthospermum australe</i>	8 seeded prostrate starbur

FARMESAFEN 250 SL will only provide variable weed control of the following weeds when sprayed at 1 ℓ /ha at seedling up to the 4 leaf stage.

<i>Bidens pilosa</i>	Black Jack
<i>Tribulus terrestris</i>	Dubbeltjie
<i>Tagetes minuta</i>	Khakiweed

Weed species that were not present during the development trials of the product could possibly also be controlled to a greater or lesser degree. The registration holder does not accept any responsibility for unlisted weeds.



## WAARSKUWINGS:

**Gevaarstellings:** Kan skadelik wees indien ingesluk. Kan skadelik wees in kontak met die vel. Veroorsaak ligte velirritasie. Kan 'n allergiese velreaksie veroorsaak.

## Onthoudingsperiodes:

Droë bone, groenbone en Sojabone..... 14 dae  
Grondbone.....56 dae

**NOTA:** Bogenoemde onthoudingsperiodes is gebaseer op plaaslike vereistes vir residuvalle en mag verskil van ander lande waarheen behandelde produkte moontlik uitgevoer kan word. Indien die behandelde gewas bestem is vir uitvoer, raadpleeg die betrokke uitvoer liggaam rakende die voor-oes interval vereistes van die lande wat die produkte uitgevoer sal word.

- **FARMESAFEN 250 SL** is getoets op die belangrikste kommersiële kultivars sonder enigefitotoksiese effekte. Meer vatbare kultivars kan in die toekoms vrygestel word wat nodig het om getoets te word. Vir dit doel dien **FARMESAFEN 250 SL** 'n beperkte toets area voordat kommersiële toepassings word gemaak.
- Hanteer versigtig
- Mag oë en vel irritasie veroorsaak.
- Berg in die oorspronklike houer in 'n veilige plek weg van voedsel en voer af.
- Hou buite bereik van kinders, oningeligte persone en diere.
- **In die geval van vergiftiging ontbied 'n genesheer en toon hulle die etiket aan.**
- **Herbetreding** - Moet nie die behandelde gebied binne 1 dag na toediening betree nie tensy beskermende oorklere gedra word.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie. Die werking en doeltreffendheid daarvan kan beïnvloed word deur faktore soos abnormale grond-, klimaats- en bergingstoestande, kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui word nie en die voorkoms van weerstand van die onkruid teen die betrokke middel, sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei en die omgewing of vir nadelige invloede op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van versuim van die gebruiker om die etiketaanwysings na te kom, of as gevolg van ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

## VOORSORGMATREËLS:

### Voorsorgstellings

**Algemeen:** Indien mediese advies benodig word, hou produkhouer of etiket byderhand. Hou buite bereik van kinders. Lees etiket voor gebruik.  
**Voorkoming:** Gebruik slegs buite of in 'n goed geventileerde area. Besoedelde werksklere moet nie uit die werkplek toegelaat word nie. Vermoed vrylating na die omgewing. Dra beskermende handskoene/beskermende klere/oogbeskerming/gesigbeskerming.  
**Reaksie:** INDIEN OP DIE VEL: Was met baie seep en water. INDIEN INGESAEM: Verwyder persoon na vars lug en hou gemaklik om asem te haal. Bel dadelik 'n dokter. INDIEN IN OË: Spoel versigtig uit met water vir 'n paar minute. Verwyder kontaklinse, indien teenwoordig en maaklik om te doen. Gaan voort met spoel. Kry mediese advies/aandag as jy onwel voel. Indien velirritasie voorkom: Kry mediese advies/aandag. Indien velirritasie of uitslag voorkom: Kry mediese advies/aandag. Trek besoedelde klere uit en was dit voor hergebruik. Versamel gemors.  
**Berging:** Berg in 'n goed geventileerde plek. Hou die houer dig toe. Stoor toegesluit.  
**Wegdoening:** Gooi inhoud en houer weg na 'n goedgekeurde afvalverwyderingsaanleg.

- Moenie rook, eet of drink terwyl die produk gehanteer word.
- Vermoed besoedeling van voedsel, voer, drinkwater asook eetgerei.
- **SPOEL die leë houer DRIEKEER UIT soos volg:** Keer die leë houer om oor die spuit- of mengtenk en dreineer vir ten minste 30 sekondes nadat die vloeit tot drup verlangsaam het. Spoel die leë houer daarna drie keer agtereenvolgens uit met een kwart van die houer volume vars water en gooi die speelwater in die spuit- of mengtenk. Prik die drievoudig uitgespoelde houer deur en gooi weg via 'n goedgekeurde versamelaar of herwinner ([www.crop-life.co.za](http://www.crop-life.co.za)). Moet nie die houer begrawe, verbrand of skenk aan enige ander partye wat dit as 'n houer vir kos of drank kan gebruik nie.
- Tydens vermenging, vermy kontak met die spuitstof so veel as moontlik.
- Maak toedieningsapparaat deeglik skoon na gebruik en gooi die waswater waar dit nie gewasse, weiding, riviere en damme sal besoedel nie.
- Vernietig die leë houer en moet dit nie vir enige ander doel gebruik nie.

## RELEVANTE STOWWE

NAAM	PRODUK IDENTIFISEERDER
Fomesafen TS	CAS-Nr.: 72178-02-0
Alkiel poligukosied	CAS-Nr.: 68515-73-1
Kalsiumdodesieibenseensulfonaat	CAS-Nr.: 26264-06-2
1,2- Bensisotiazolin-3-een	CAS-Nr.: 2634-33-5

## SIMPTOME VAN VERGIFTIGING

Hoofpyn, duiseligheid, naarheid.

## NOODHULPBEHANDELING

Moenie braking veroorsaak nie. Kry mediese hulp.

## NOTA AAN GENEESHEER

Simptomatiese behandeling.

## WEERSTANDSWAARSKUWING

Vir weerstands bestuur, is **FARMESAFEN 250 SL** 'n groep kode 14 onkruidodder. Enige populasie van 'n spesifieke onkruid mag individuele onkruide insluit wat 'n natuurlike weerstand teen **FARMESAFEN 250 SL** en ander groepkode 14 onkruidodders het. Indien hierdie onkruidodders herhaaldelik aangewend word, kan die weerstandbiedende individue uiteindelik die onkruidpopulasie oorheers Hierdie weerstandbiedende onkruide sal waarskynlik nie deur **FARMESAFEN 250 SL** of enige ander groepkode 14 onkruidodder beheer word nie.

## Om weerstand teen onkruidodders te vertraag:

- Vermoed die eksklusiewe en herhaaldelike gebruik van onkruidodders met dieselfde groepkode. Wissel af met, of gebruik tenkmengsels van, produkte in verskillende onkruidodder groepkodes.
- Integreer ander beheermaatreëls (chemies, meganies, biologies) in onkruidodder programme.

Vir spesifieke inligting oor weerstandsbestuur raadpleeg die registrasiehouer van hierdie produk.

Die teenwoordigheid van sulke weerstandbiedende onkruide is moeilik om te bepaal voordat die onkruidodder toegedien word. Dus kan die registrasiehouer en die verspreider nie verantwoordelikheid aanvaar vir enige verlies wat mag voortspruit uit **FARMESAFEN 250 SL** se nie weerstandbiedende onkruid beheer. Moenie **FARMESAFEN 250 SL** alleen in situasies wat weerstand bevestig is, toedien nie. Sub- standaard toedienings van **FARMESAFEN 250 SL** sal lei tot onbevredigende onkruidbeheer wat die ontwikkeling van onkruidodderweerstand kan vorder.

## WYSE VAN AKSIE

Protoporfirinogeen oksidase inhibeerder. Selektiewe onkruidodder, geabsorbeer deur beide blare en wortels, met baie beperkte translokasie in die floëem.

## GBRUIKSBEPERKINGS:

1. **FARMESAFEN 250 SL** moet nie gebruik word waar sojabone, droëbone, groenbone of grondbone tussen ander gewasse geplant is.
2. Alle ander gewasse, behalwe vir die wat op die produk geregistreer is, moenie binne 12 maande van **FARMESAFEN 250 SL** toediening, geplant word nie.
3. Moenie **FARMESAFEN 250 SL** toedien op enige gewas of situasie wat nie op hierdie etiket verskyn is nie.
4. **FARMESAFEN 250 SL** moet nie in die Noordwes-provinsie of die Wes-Vrystaat gebruik word nie.
5. **FARMESAFEN 250 SL** kan slegs op die volgende kultivars gebruik word:
  - Sojabone - alle bestaande kultivars.
  - Droëbone - Nuweveld, Teebus en Bonus, World Cup, ORS
  - Groenbone - Strike, Provider, Cape, Derby, Palati, Newton
  - Gondbone - Sellie.
6. Moenie hierdie produk in kombinasie of in tenkmengsels toedien met enige ander produkte of landboumiddels wat nie spesifiek op die etiket gelys is vir toediening nie. Cedar Falls Properties 130 (Pty) Ltd aanvaar geen verantwoordelikheid vir enige negatiewe gevolge as die instruksies op die etiket nie nagekom word nie, tensy 'n Meridian verteenwoordiger geraadpleeg is.



## **GEBRUIKSAANWYSINGS    Gebruik slegs soos aangedui**

**FARMESAFEN 250 SL** is 'n selektiewe na-opkoms breëblaardoder wat tot die voor-blom stadium van die gewas teodien word, **maar moet nie later as 4 weke na plant en nie later as die 4- blaarstadium van die onkruid teodien word nie.**

Toediening op 'n later groeistadium kan die kans van 'n paar weergroei van behandelde onkruid verhoog. Dit word aanbeveel om te wag totdat die meerderheid van breëblaar onkruid ontkiem het voor bespuiting met **FARMESAFEN 250 SL** (verwys na die onkruid lys hieronder). Reën vanaf 4 - 6 uur na toediening sal geen invloed op die aktiwiteit van **FARMESAFEN 250 SL** hê nie.

Indien die normale verbouing metode is die gebruik van voor-opkoms onkruidodders, dan moet hierdie praktyk voortgesit word gevolg deur **FARMESAFEN 250 SL** vir later onkruid ontkieming.

Tydlike simptome van gewasskade , sirkulere brandmerkie op die blare en die moontlike vertraging van groei mag voorkom na 'n **FARMESAFEN 250 SL** toediening, maar dit sal nie tot oesverlies lei nie.

## **VERENIGBAARHEID**

Moet nie **FARMESAFEN 250 SL** met ander produkte meng nie.

**FARMESAFEN 250 SL** mag in program met ander onkruidodders gebruik word voorsienende dat genoegsame tyd intervalle toegelaat word sodat ander onkruid (soos grasse) nie onder stress geplaas word nie. Byvoorbeeld, waar **FARMESAFEN 250 SL** voor of nadat 'n toediening van fluaizifop-p-butyl bevattende produkte gebruik word, moet 14 dae tussen bespuiting plaasvind.

Wanneer die gebruik van ander produkte in 'n spuitprogram met **FARMESAFEN 250 SL** plaasvind, moet die etikette van die spesifieke produkte met betrekking tot (1) aanwysings vir hul toediening en (2) wisselbou aanbevelings altyd geraadpleeg word.

## **KLIMAATSTOESTANDE:**

**FARMESAFEN 250 SL** moet toegedien word wanneer die onkruid aktief groei onder gunstige temperatuur en vog toestande. **FARMESAFEN 250 SL** moet nie toegedien word wanneer die gewasse en onkruid onder droogtestremming of koue temperature is of wanneer swak onkruidbeheer tot gevolg sal wees nie.

## **GRONDGEDRAAGDE SIEKTES**

Moenie **FARMESAFEN 250 SL** op sojabone, groenbone, grondbone of droëbone wat reeds simptome van grond siektes toon toedien nie omdat die kombinasie van 'n **FARMESAFEN 250 SL** behandeling en siektestres kan lei tot stand verliese.

## **TOEDIENINGSMETODES**

Verseker dat die toerusting korrek gekalibreer is en gereeld nagegaan word tydens toediening om 'n egalige en akkurate toediening te verseker. As reën verwag word binne 4-6 ure, moet nie **FARMESAFEN 250 SL** toedien nie. **FARMESAFEN 250 SL** kan deur grondtoerustings toegedien word. Gebruik slegs skoon water.

## **GRONDTOEDIENING**

Dien **FARMESAFEN 250 SL** toe met minimum van 200 - 300 L water / ha. Gebruik platwaaier-of holkeëlsputpunte met 'n spuitdruk van 150kPa. Verseker dat die spuitmengsel eweredig goeie bedekking van die onkruid sal gee.

## **AANBEVELINGS**

<b>GEWAS</b>	<b>FARMESAFEN 250 SL ℓ /ha</b>	<b>OPMERKINGS</b>
<b>Sojabone, Droëbone, Groenbone, Grondbone.</b> Slegs breëblaaronkruid	1 ℓ / ha	Spuit wanneer die onkruid tussen die saailinge en 4-blaarstadium is en voor blom van die gewas. Verwys hieronder vir die onkruid wat normalweg beheer sal word.

**ONKRUIDE WAT NORMAALWEG BEHEER SAL WORD:**

Die volgende breëblaaronkruid (saailinge tot en met die 4 blaar stadium) sal deur

**FARMESAFEN 250 SL op 1 ℓ / ha beheer word:**

<i>Amaranthus hybridus</i>	Kaapse misbredie
<i>Datura ferox</i>	Grootstinkblaar
<i>Galinsoga parviflora</i>	Knopkruid
<i>Portulaca oleracea</i>	Porslein
<i>Schkuhria pinnata</i>	Kleinkakiebos
<i>Sida cordifolia</i>	Hartblaartaaiman
<i>Richardia brasiliensis</i>	Tropiese Richardia
<i>Acanthospermum australe</i>	Agtsadige kruipsterklits

**FARMESAFEN 250 SL** sal slags wisselvallige onkruid beheer van die volgende onkruid wanner dit op 1 ℓ / ha op saailinge tot en met die 4 blaar stadium bespuit word.

<i>Bidens pilosa</i>	Black Jack
<i>Tribulus terrestris</i>	Dubbeltjie
<i>Tagetes minuta</i>	Khakiweed

Ander onkruidspesies wat nie tydens die ontwikkelingsproewe van die produk voorgekom het nie, kan moontlik in 'n mindere of meerdere mate beheer word. Die registrasiehouer aanvaar egter geen aanspreeklikheid ten opsigte van ongelyste onkruid nie.



# FARMESAFEN 250 SL

Reg. No. / Nr L10319 Act No. 36 of 1947/ Wet Nr 36 van 1947

HERBICIDE



**READ THE LABEL AND ATTACHED PACKAGED LEAFLET BEFORE USE  
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

HRAC HERBICIDE GROUP CODE

14

ONKRUIDDODER GROEPKODE

A post-emergence water soluble concentrate selective herbicide for the selective control of various annual broadleaf weeds in the crops as listed.

'n Na-opkoms water oplosbare konsentraat selektiewe onkruiddoder vir die selektiewe beheer van verskeie eenjarige breëblaaronkruid in gewasse soos aangedui.

## ACTIVE INGREDIENT/AKTIEWE BESTANDEEL:

Fomesafen..... 250 g/ ℓ



**DANGER**

### Hazard Statements:

May be harmful if swallowed.  
May be harmful in contact with skin.  
Causes mild skin irritation.  
May cause an allergic skin reaction.  
Causes serious eye damage.  
Toxic if inhaled.  
May cause damage to organs through prolonged or repeated exposure.  
Very toxic to aquatic life with long lasting effects.

### Precautionary Statements:

Do not breathe spray, vapours.  
IF ON SKIN: Wash with plenty of soap and water.  
IF INHALED: Remove person to fresh air and keep comfortable for breathing. Immediately call a doctor.  
IF IN EYES: Rinse cautiously with water for several minutes. Immediately call a doctor.  
If skin irritation or rash occurs: Get medical advice/attention.

Batch No:

D.O.M:

D.O.E:

Net Mass:

See reverse of packaging for details.

FOR FULL PARTICULARS SEE ATTACHED PAMPHLET.  
VIR VOLLEDIGE BESONDERHEDE KYK AANGEHEGTE ETIKET.

UN No. / VR Nr: 3082

Poison Information Helpline (24 hrs): 0861 555 777



REGISTRATION HOLDER/ REGISTRASIEHOUER  
Cedar Falls Properties 130 (Pty) Ltd  
Reg. Nr.: Reg. No. 2006/031383/07  
61 Marshall Drive, Old Mill Industrial Park,  
Mount Edgecombe, South Africa, 4300  
Tel: 031 003 3486



BARCODE

100491

165(H) x 140(W) mm

CYAN

MAG

YEL

BLK

100491