

WARNINGS

Hazard Statements: : Fatal if swallowed or if inhaled. May be harmful in contact with skin. Harmful to aquatic life with long lasting effects.

PRE-HARVEST INTERVAL: The following minimum number of days must be allowed between last application and harvest:

Table potatoes.....40 days
Sugarcane.....120 days
Tomatoes.....21days

Handle with extreme care:

- Toxic to fish, bees and wildlife.
- Keep out of reach of children, uninformed persons and animals.
- **Re-entry Interval:** In case of re-entry of treated area within 30 days of application or handling treated soil within 30 days of application, protective clothing (overalls, rubber boots and gloves) should be worn.
- In case of poisoning, immediately call a doctor and make this label available to him/her.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions that could not have been foreseen in terms of the registration. Consult the Supplier in the event of any uncertainty.

PRECAUTIONS

Precautionary Statements

General: If medical advice is needed, have product container or label at hand. Keep out of reach of children. Read label before use.

Prevention: Do not breathe dust. Wash hands, forearms and face thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection. In case of inadequate ventilation wear respiratory protection.

Response IF SWALLOWED: Immediately call a poison centre or doctor. IF INHALED: Remove person to fresh air and keep comfortable for breathing. Immediately call a poison centre or doctor. IF exposed or concerned: Get medical advice/attention. Call a doctor if you feel unwell. Get medical advice/attention if you feel unwell. Specific treatment is urgent (see ANTIDOTE, supplemental first aid instruction on this label).

Storage: Store in a well-ventilated place. Keep container tightly closed. Store locked up

Disposal: Dispose of contents and container to an approved waste disposal plant.

- **CLEAN THE EMPTY FOIL-LINED OR PLASTIC-LINED PAPER BAG AS FOLLOWS:** Shake all the pesticide formulation out of the bag into the spray or mixing tank. Turn the bag inside out. Hold the bag over the spray or mixing tank and rinse the bag with running water for thirty seconds. Shake the rinsate off into the tank. Repeat the rinsing twice. Do not bury, burn or donate the bag to any other parties that may use it as a container for foodstuffs. These bags cannot be recycled and must be incinerated by a licensed incineration facility.
- Prevent contamination of foods, feedstuffs, drinking water and eating utensils.
- Clean applicator after use, disposing of wash water where it will not contaminate crops grazing, rivers and dams.

RELEVANT SUBSTANCES

NAME	PRODUCT IDENTIFIER
Silica sand	CAS-No.: 14808-60-7
Oxamyl TC	CAS-No.: 23135-22-0
Calcium lignosulfonate	CAS-No.: 8061-52-7

SYMPTOMS OF HUMAN POISONING Headache, fatigue, faintness, giddiness, excessive sweating, nausea, abdominal pains, vomiting, blurred vision, muscle twitching, unusually small pupils, respiratory distress, coma.

RESTRICTED USE AGRICULTURAL REMEDY

THIS REMEDY IS RESTRICTED DUE TO ACUTE TOXICITY. THIS REMEDY MAY ONLY BE SOLD TO AND USED BY A REGISTERED PEST CONTROL OPERATOR, OR BY SOMEONE UNDER THE SUPERVISION OF A REGISTERED PEST CONTROL OPERATOR, AND ONLY FOR THOSE USES COVERED BY THE PEST CONTROL OPERATOR'S SCOPE OF REGISTRATION, AND ONLY AS DIRECTED ON THIS LABEL.

FOXAMYL 10G

Reg. No. / Nr L9877 Act No. 36 of 1947/ Wet Nr 36 van 1947

NEMATICIDE



**READ THE LABEL AND ATTACHED PACKAGED LEAFLET BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

INSECTICIDE / NEMATICIDE GROUP CODE

1A

ONKRUIJDODER GROEPKODE

A granular remedy for protection of potatoes and tomatoes against root knot nematodes (*Meloidogyne spp.*) and the control of nematodes in sugar cane growing in Kwa-Zulu Natal.

'n Middel in korrelvorm vir beskerming van aartappels en tamaties teen knopwortelaalwurm (*Meloidogyne spp.*), asook die beheer van aalwurm in suikerriet wat in Kwa-Zulu Natal groei.

ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL:

Oxamyl (carbamate)100 g/kg..... Oksamil (karbamaat)



DANGER

Batch No:

D.O.M:

D.O.E:

Net Contents:

See reverse of packaging for details.

FOR FULL PARTICULARS SEE ATTACHED PAMPHLET.
VIR VOLLEDIGE BESONDERHEDE KYK AANGEHEGTE ETIKET.

UN No. / VR Nr: 2757



REGISTRATION HOLDER/ REGISTRASIEHOUER
Cedar Falls Properties 130 (Pty) Ltd
Reg. Nr: Reg. No. 2006/031383/07
61 Marshall Drive, Old Mill Industrial Park,
Mount Edgecombe, South Africa, 4300
Tel: 031 003 3486



Scan for GHS
and label details

BARCODE

100512

FIRST AID TREATMENT

Remove patient from source of poisoning to a cool and well-ventilated area. Calm him and keep him still. Remove contaminated clothing and rinse contaminated body area thoroughly with plenty of soap and cold water. Do not rub skin. Flush contamination from eyes with clean water for at least 15 minutes. If swallowed, induce vomiting by tickling the back part of the throat. Repeat until vomit is clear and free from the smell of poison. Do not apply direct mouth-to-mouth resuscitation. Never give anything by mouth to an unconscious person. Immediately take the patient to the nearest physician.

NOTE TO PHYSICIAN

Administer Atropine sulphate intravenously (2 - 4 mg/adult) every 10 minutes until signs of atropinization (dry flushed skin and tachycardia) appear. Maintain atropinization until the patient recovers. Do not apply Pralidoxime (2 PAM, Protopam) and/or other oximes. Do not use morphine.

RESISTANCE WARNING STATEMENT

FOXAMYL 10G is an insecticide/nematicide GROUP CODE 1A product. Any nematode population may contain individuals naturally resistant to **FOXAMYL 10G** and other GROUP CODE 1A insecticides/nematicides. The resistant individuals can eventually dominate the population if these products are used repeatedly and will then not be controlled by **FOXAMYL 10G** or by any other GROUP CODE 1A product.

To delay the development of resistance:

- Avoid exclusive repeated use of GROUP CODE 1A products. Alternate with products from other GROUP CODES.
- Integrate other control methods (e.g. crop rotation with non-susceptible crops, fallow lands etc.) into nematode control programs.

Since the occurrence of resistance is difficult to detect prior to use, it is of the utmost importance that all of the above precautions are strictly adhered to. The Registration Holder or Distributor cannot be held responsible if **FOXAMYL 10G** fails to control resistant nematodes. For further advice on the prevention of resistance, contact your Distributor's local representative.

MODE OF ACTION:

Cholinesterase inhibitor. Contact and systemic insecticide, acaricide and nematicide. Absorbed by the foliage and the roots, with translocation.

USE RESTRICTIONS

Cultural practices: Monoculture and/or crop rotation with nematode susceptible crops must be avoided.

Soil type and moisture: Light sandy soils and/or dry soil during application can result in poor control.

Soil pH: High soil pH (alkaline soil) will significantly reduce **FOXAMYL 10G**'s efficacy. Do not use **FOXAMYL 10G** on soils with a pH above 7,5 or if the pH of the irrigation water is above 7,5 without first consulting the distributor of this product or the registration holder.

Infected soil and/or high risk areas: Apply a full control programme consisting of soil fumigation followed by **FOXAMYL 10G** at planting followed by one or more foliar sprays on soils with a history of high nematode infestations and/or in high risk areas such as Barkley-West, Christiana, Coetzenradam, Douglas, Hopetown, PK le Roux Dam, Priessa, Tosca, Vaalharts and Vryburg.

Resistant nematodes: Columbia root knot nematode (*Meloidogyne chitwoodi* and *M. fallax*) are not controlled. These 2 species occur in colder areas such as Clooclane, Maclear and Ugie as well as in warmer areas such as Boston, Mooi River, Swaziland and Vaalharts. Golden cyst nematode (*Globodera pallida* and *G. rostochiensis*) is also not controlled.

Potatoes: Plant only tubers that are certified nematode free. Use only on table potatoes. Do not use on seed or chips potatoes. Do not store potatoes in the soil: lift as soon as possible after last foliar application and not later than 4 weeks.

RESTRICTED USES: Use only as directed

CROP AND PEST	DOSAGE	REMARKS
POTATOES Root knot nematodes: <i>Meloidogyne</i> species (excluding <i>Meloidogyne chitwoodi</i> and <i>M. fallax</i>)	250 g/100 m furrow	Table Potatoes only: Soil Application: Apply only to moist soil. Apply into planting furrow before closing. Foliar Application: Follow up with a foliar application of Foxamyl 310 SL and consult the registered labels for directions for use.
SUGAR CANE: For use in KwaZulu-Natal only Nematodes: Including Root knot nematodes (<i>Meloidogyne</i> species) and Lesion nematodes (<i>Pratylenchus</i> species)	30 kg/ha for 1.2 m rows or 360 g/100 m planting furrow	Plant cane only: Place granules in the furrow with the seed at planting using a mechanical granular applicator. Cover both with soil. Note: 1. FOXAMYL 10G should NEVER be applied together with lime in the plant furrow. Instead it is recommended that the lime should be broadcast during soil preparation. If lime is to be applied in the furrow, band apply FOXAMYL 10G on the soil surface alongside or over rows SOON AFTER LEAF EMERGENCE. Thoroughly incorporate granules. 2. Apply only on soils with LESS than 6% clay.
TOMATOES Root knot nematodes: <i>Meloidogyne</i> species (excluding <i>Meloidogyne chitwoodi</i> and <i>M. fallax</i>)	125 g/100 m planting furrow	Soil application: Apply only to moist soil. Apply either in the planting furrow or in a 20 cm band over the planting furrow and incorporate into the soil. Foliar Application: Follow up with a foliar application of Foxamyl 310 SL and consult the individual labels for directions for use.



WAARSUWINGS:

Gevaarstellings: Dodelik indien ingesluk of ingeasem word. Kan skadelik wees in kontak met die vel. Skadelik vir waterlewé met langdurige effekte.

ONTHOUDENGSPERIODE: Die volgende onthoudingsperiodes moet toegelaat word tussen laaste toediening en oes van die volgende gewasse:

Tafelaartappels 40 dae

Suikerriet 120 dae

Tomaties 21 dae

Hanteer uiters versigtig:

- Giftig vir visse, bye en wild.
- Giftig deur velkontak, innname deur mond en inaseming.
- **Herbetreding:** Indien die land binne 30 dae na behandeling betree of die grond binne 30 dae na behandeling hanteer word moet beskermende klerk (oorkap, rubberstewels en handskoene) gedra word.
- In geval van vergiftiging, onttrek onmiddellik 'n geneesheer en ton hierdie etiket aan hom/haar.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie, aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale grond-, klimaats- en bergingstoestande, verenigbaarheid met ander stowwe wat nie op die etiket aangedui word nie en die voorkoms van weerstand van die plaag teen die betrokke middel, sowel as die metode, tyd en akkurateheid van toediening. Verder aanvaar die registrasiehouer geen verantwoordelikheid vir skade aan gewasse, plantgroei, die omgewing of vir die nadelige effek op mens of dier of vir 'n gebrek aan prestaties as gevolg van die versuum van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die Verspreider in geval van enige onsekerheid.

VOORSORGMAATREËLS

Voorsorgstellings

Algemeen: Indien mediese advies benodig word, hou produkhouer of etiket byderhand. Hou buite bereik van kinders. Lees etiket voor gebruik. **Voorkoming:** Moet nie stof inasem nie. Was hande, voorarms en gesig deeglik na hantering. Moet nie eet, drink of rook wanneer hierdie produkt gebruik word nie. Gebruik slegs buite of in 'n goed geventileerde area. Verminder vrylating na die omgewing. Dra beskermende handskoene/ beskermende klere/oogbeskerming/gesigbeskerming. Dra asemhalingsbeskerming in geval van onvoldoende ventilasie.

Reaksie: INDIEN INGESLUK: Ontbied onmiddellik 'n gifsentrum of dokter. INDIEN INGESAEM: Verwyder persoon na vars lug en hou gemaklik om asem te haal. Bel dadelik 'n gifsentrum of dokter. INDIEN BLOOTGESTEL OF BEKOMMERD: Kry mediese advies/aandag. Bel 'n dokter as jy onwel voel. Kry mediese advies/aandag as jy onwel voel. Spesifieke behandeling is dringend (sien teenmiddel, aanvullende noodhulpinstruksies op hierdie etiket).

Berging: Berg in 'n goed geventileerde plek. Hou die houer dig toe. Stoor toegesluit.

Wegdoening: Gooi inhoud en houer weg na 'n goedgekoonde afvalverwyderingsaanleg.

- **MAAK DIE LEË FOELIE-GEVOERDE OF PLASTIEK-GEVOERDE PAPIERSAK SKOON SOOS VOLG:** Skud al die plaagdoderformulerig uit die sak in die spuit- of mengtenk. Draai die sak binnekste na buite. Hou die sak oor die spuit- of mengtenk en spoel die sak uit met lopende water vir dertig sekondes. Skud die spoelwater in die tenk af. Herhaal die spoel twee keer. Moet nie die sak begrawe, verbrand of skenk aan enige ander partye wat dit as 'n houer vir voedsel kan gebruik nie. Hierdie sakke kan nie herwin word nie en moet deur 'n gelisensierte verbrandingsfasilitet verbrand word.
- Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.
- Maak die toedieningsapparaat deeglik skoon na gebruik. Verwyder uitspoelwater na waar dit nie ander gewasse, weiding, riviere en damme sal besoedel nie.

RELEVANTE STOWWE

NAAM	PRODUK IDENTIFISEERDER
Silika sand	CAS-Nr.: 14808-60-7
Oksamiel TS	CAS-Nr.: 23135-22-0
Kalsium lignosulfonaat	CAS-Nr.: 8061-52-7

SIMPTOME VAN VERGIFTIGING

Hooypyn, vermoedelheid, flouheid, duiselheid, oormatige sweat, naarheid, maagpyn, braking, versteurde gesigsvermoë, spiertrekings, buitengewoon verkleinde pupille, asemhalingsnood en bewusteloosheid.

NOODHULPBEHANDELING

Verwyder die persoon van die oorsaak van die vergiftiging na 'n koel en goed geventileerde omgewing. Kalmeer hom en hou hom stil. Besmette oë moet vir minstens 15 minute met skoon water uitgespoel word. Verwyder besoedelde klere en was die besmette liggaaumsdele met seep en koue water. Moenie die vel hard vryf nie. As die produk per mond ingeneem is, veroorsaak braking deur die agterste gedeelte van die keel met u vinger te prikkel. Herhaal tot braaksel helder is en nie meer na gif ruik nie. Moenie direkte mond tot mond asemhaling toepas nie. Moet nooit iets per mond aan 'n bewusteloze persoon gee nie. Neem die persoon onmiddellik na die naaste geneesheer en stel hierdie etiket aan hom/haar beskikbaar.

NOTA AAN GENEESHEER

Dien atropensiulfat binneaars elke 10 minute toe (2 – 4 mg/volwassene) tot die eerste tekens van atropinisasie (droë, gloeiende vel en tagkardie) voorkom. Hou aan met atropinisasie tot die pasiënt herstel. Moenie Praliedoksieme (2-PAM, Protopam) of ander oksiemes in plaas van atropine toedien nie. Moenie morfiën gebruik nie.

VOORKOMING VAN WEERSTAND EN BESTUUR

FOXAMYL 10G is 'n aalwurm/insekddoder GROEPKODE 1A middel. Enige aalwurm populasie kan individue bevat wat natuurlik bestand is teen **FOXAMYL 10G** en ander GROEPKODE 1A middels. Hierdie bestande individue kan uiteindelik die populasie oorheers indien hierdie middels herhaaldelik gebruik word en sal dan nie deur **FOXAMYL 10G** of ander GROEPKODE 1A middels beheer word nie.

Om die voorkoms van weerstand te vertraag:

- Vermy die herhaaldelike gebruik van GROEPKODE 1A middels. Wissel af met middels van ander GROEPKODES.
- Integreer ander beheermaatreëls (bv. wisselbou met nie vatbare gewasse, braakké ens) in aalwurmbeheer programme.

Aangesien die voorkoms van weerstand moeilik is om te bepaal voor behandeling is dit uiters belangrik dat al die bogenoemde voorsorgmaatreëls streng nagekom word. Die Registrasiehouer van Verspreider kan nie verantwoordelik gehou word indien **FOXAMYL 10G** nie weerstandsbiedende aalwurms beheer nie. Vir verdere advies m.b.t. die voorkoming van weerstand, kontak u plaaslike verteenwoordiger van die Verspreider.

WYSE VAN AKSIE

Cholinesterase inhibeerder. Kontak- en sistemiese insekdoder, mytdoder en aalwurmdoder. Geabsorbeer deur die blare en die wortels, met translokasie.

GEBRUIKSBEPERKINGS

Verbouingspraktyke: Monokultuur en/of wisselbou met aalwurm vatbare gewasse moet vermy word.

Grondtipe en -vog: Ligte/sanderige grond en/of droë grond tydens toediening kan tot onvoldoende beheer lei.

Grond pH: Höe grond-pH (m.a.w. alkaliese grond) sal **FOXAMYL 10G** se effektiwiteit negatief beïnvloed. Moenie **FOXAMYL 10G** gebruik op grond met 'n pH bo 7,5 of indien die pH van die besproeiingswater bo 7,5 is nie. Raadpleeg eers die verspreider of die registrasiehouer van die produk.

Besmette grond en/of hoë risiko areas: Dien 'n volledige beheerprogram toe nl. 'n vooraf grondberoking gevvolg deur **FOXAMYL 10G** tydens plant gevvolg deur een of meer blaarbespuitings op lande met 'n geskiedenis van hoë aalwurm besmetting en/of in hoë risiko areas soos Barkley- Wes, Christiana, Coetzersdam, Douglas, Hopetown, PK le Roux Dam, Prieska, Tosca, Vaalharts en Vryburg.

Bestande aalwurms: Columbia knopwortel aalwurm (*Melodogyne chitwoodi* en *M. fallax*) word nie beheer nie. Hierdie 2 spesies kom voor in kouer gebiede soos Clocolane, Maclear en Ugie asook in warmer dele soos Boston, Moorivier, Swaziland en Vaalharts. Goue sist aalwurm (*Globodera pallida* en *G. rostochiensis*) word ook nie beheer nie.

Aartappels: Plant slegs gesertifiseerde aalwurmvrye knolle. Gebruik slegs op tafelaartappels: nie op saad- of skyfie aartappels nie. Moenie aartappels in die grond stoor nie: haal so gou as moontlik uit na laaste blaarbespuiting en nie later as 4 weke nie.

BEPERKTE GEBRUIK: Gebruik slegs soos aangedui

GEWAS EN PLAAG	DOSIS	OPMERKINGS
AARTAPPELS Knopwortelaalwurms Meloidogyne spesies (<i>uitsluitend</i> Meloidogyne chitwoodi en M. fallax)	250 g/100 m plantry	Slegs tafelaartappels: Grondtoediening: Dien slegs in klam grond toe. Dien in die plantvoor toe tydens plant voordat dit toegemaak word. Blaarbespuitings: Volg op met 'n blaarbespuiting van Foxamyl 310 SL en raadpleeg die geregistreerde etikette vir gebruiksaanwysings.
SUIKERRIET For use in KwaZulu-Natal only Aalwurms Insluitend knopwortelaalwurms (Meloidogyne spesies) en MeloidogyneLesion aalwurms (<i>Pratylenchus</i> spesies)	30 kg/ha vir 1.2 m ry spasiëring of 360 g/100 m plantvoorlengte	Slegs plant-oes: Plaas korrels met 'n meganiese korreltoedienier in die plantvoor by steggies. Bedek beide met grond. Nota: 1. MOENIE FOXAMYL 10G saam met kalk in die plantvoor toedien nie. Kalk moet verkiekslik breedwerpig tydens grondvoorbereiding toegedien word, Waar kalk wel in die plantvoor toegedien is moet FOXAMYL 10G op die grondoppervlak in 'n strook bo-oor of aan weerskante van die ry toegedien word SODRA DIE BLARE VERSKYN en ry sigbaar is. Werk die korrels deeglik in. 2. Gebruik slegs op grond met 'n klei-inhoud van minder as 6%.
TAMATIES Knopwortelaalwurms Meloidogyne spesies (excluding Meloidogyne chitwoodi en M. fallax)	125 g/100 m plantry	Grondtoediening: Dien slegs in klam grond toe. Dien óf in die plantry, óf in 'n 20 cm strook óór die plantry toe, en werk in die grond in. Blaarbespuitings: Volg op met 'n blaarbespuiting van Foxamyl 310 SL en raadpleeg die geregistreerde etikette vir gebruiksaanwysings.



RESTRICTED USE AGRICULTURAL REMEDY

THIS REMEDY IS RESTRICTED DUE TO ACUTE TOXICITY. THIS REMEDY MAY ONLY BE SOLD TO AND USED BY A REGISTERED PEST CONTROL OPERATOR, OR BY SOMEONE UNDER THE SUPERVISION OF A REGISTERED PEST CONTROL OPERATOR, AND ONLY FOR THOSE USES COVERED BY THE PEST CONTROL OPERATOR'S SCOPE OF REGISTRATION, AND ONLY AS DIRECTED ON THIS LABEL.

FOXAMYL 10G

Reg. No. / Nr L9877 Act No. 36 of 1947/ Wet Nr 36 van 1947

NEMATICIDE



**READ THE LABEL AND ATTACHED PACKAGED LEAFLET BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

INSECTICIDE / NEMATICIDE GROUP CODE

1A

ONKRUIJDODER GROEPKODE

A granular remedy for protection of potatoes and tomatoes against root knot nematodes (*Meloidogyne spp.*) and the control of nematodes in sugar cane growing in Kwa-Zulu Natal.

'n Middel in korrelvorm vir beskerming van aartappels en tamaties teen knopwortelaalwurm (*Meloidogyne spp.*), asook die beheer van aalwurm in suikerriet wat in Kwa-Zulu Natal groei.

ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL:

Oxamyl (carbamate)100 g/kg..... Oksamil (karbamaat)



DANGER

Batch No:

D.O.M:

D.O.E:

Net Contents:

See reverse of packaging for details.

FOR FULL PARTICULARS SEE ATTACHED PAMPHLET.
VIR VOLLEDIGE BESONDERHEDE KYK AANGEHEGTE ETIKET.

UN No. / VR Nr: 2757



REGISTRATION HOLDER/ REGISTRASIEHOOUER
Cedar Falls Properties 130 (Pty) Ltd
Reg. Nr: Reg. No. 2006/031383/07
61 Marshall Drive, Old Mill Industrial Park,
Mount Edgecombe, South Africa, 4300
Tel: 031 003 3486



Scan for GHS
and label details

BARCODE

100512

165(H) x 140(W) mm

CYAN MAG YEL BLK

100512