

WARNINGS

Withholding periods: (Minimum number of days between last application and harvest or feeding / grazing).

Table grapes	Not later than pre-blossom stage.
---------------------	--

- Handle with care.
- Eye irritant and possible skin sensitizer upon contact.
- Toxic to fish.
- Store in a cool, dry place away from food, feeds, fertilizers and other agricultural remedies.
- Keep out of reach of children, animals and uninformed persons.
- Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be effective under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment, or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

- Avoid eye and skin contact and inhalation of spray mist.
- Wear rubber gloves and a face shield when handling the concentrate and application of the spray mixture.
- In case of eye contact, flush out at once.
- In case of skin contact, wash off at once.
- In case of spillage of product on clothes, change and wash immediately.
- Do not eat, drink or smoke whilst mixing, applying the product or before washing hands and face or change of clothing.
- Avoid drift of spray onto other crops, grazing, rivers, dams and areas not under treatment.
- Thoroughly clean the spraying equipment directly after use and dispose of wash water where it will not contaminate food, grazing, boreholes, rivers or dams.
- Invert the empty container over the spray or mixing tank to make sure the container is empty before destroying the container.
- **TRIPLE RINSE** empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of 10 % of that of the container. Add the rinsing to the contents of the spray tank before destroying the container in the prescribed manner.
- Destroy the empty container by perforation and flattening.
- **Never** re-use the empty container for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

FIRST AID TREATMENT:

IF IN EYES:	Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Remove contact lenses. Immediate medical attention is required.
IF SWALLOWED:	Rinse mouth and then drink plenty of water. Do not induce vomiting, unless told by a poison control centre, or doctor. Seek medical attention. Never induce vomiting, or give anything by mouth, if the victim is unconscious, or having convulsions.
IF ON SKIN:	Take off all contaminated clothing immediately. Wash off immediately with plenty of water. If skin irritation persists, call a physician. Wash contaminated clothing before re-use.
IF INHALED:	Remove to fresh air. If breathing is irregular or stopped, administer artificial respiration. Keep patient warm and at rest. Call a physician or Poison Information Centre immediately.

RESISTANCE WARNING:

FARM-AG FOLPET 500 SC is a group code M4 fungicide. Any fungus population may contain individuals naturally resistant to **FARM-AG FOLPET 500 SC** and other group code M4 fungicides. The resistant individuals can eventually dominate the fungus population if these fungicides are used repeatedly and exclusively in programs. These resistant fungi may not be controlled by **FARM-AG FOLPET 500 SC** or any other group



FARM-AG FOLPET 500 SC

Reg No/ Nr L10591 Act No/ Wet Nr 36/1947

A suspension concentrate organic based fungicide for the preventive control of deadarm disease, anthracnose and downy mildew preblossom in table grapes as listed.

'n Suspensie konsentraat organiesgebaseerde swamdoder vir die voorkomende beheer van streeppleksiwkte, antraknose en donsige skimmel voor-blom in tafeldruwe.

FRAC FUNGICIDE GROUP CODE

M4

FRAC SWAMDODER GROEPKODE

ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL:

Folpet (phtalimide)..... 500 g/l..... Folpet (ptalimied)

Manufacture Date:

Vervaardigingsdatum:

Batch No.:

Lotnommer:

NET MASS/ INHOUD

l

**SEE ENCLOSED LEAFLET FOR FULL PARTICULARS
SIEN INGESLOTE VOUBILJET VIR VOLLE BESONDERHEDE**

REGISTRATION HOLDER/REGISTRASIEHOUER:

Cedar Falls Properties 130 (Pty) Ltd.

Co. Reg.No. 2006/031383/07

61 Marshall Drive, Old Mill Industrial Park, Mt Edgecombe, South Africa, 4300 • Tel. (031) 003 3486

Emergency Number: Griffon Poison Information Centre

UN No.: 3082



**CAUTION
VERSIGTIG**



code M4 fungicides.

To delay fungicide resistance:

- Avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate with products from different fungicide group codes,
- Integrate other control methods (chemical, cultural, biological) into insect control programmes.

For specific information on resistance management contact the registration holder of this product.

Compatibility:

FARM-AG FOLPET 500 SC can be applied in a tank-mix with most of the commonly used insecticides and fungicides with the exception of Lime, Lime Sulphur and Bordeaux mixture.

DIRECTIONS FOR USE: Use only as directed

Compatibility:

- **FARM-AG FOLPET 500 SC** can be applied in tank mixtures with the most commonly used insecticides and fungicides with the exception of Lime, Lime Sulphur and Bordeaux mixture. Do not apply more than four applications per season.
- It should not be applied with any other Botrytis remedy table grapes.

Mixing instructions:

Shake the container thoroughly before tank mixture is prepared. Half fill the spray tank with water and agitate. After the pre-mix tank is half full with water, add **FARM-AG FOLPET 500 SC** and top up with water to the required volume while agitating continuously. Do not use brackish water. Re-suspend the spray mixture if agitation was suspended for longer than 5 minutes. If **FARM-AG FOLPET 500 SC** is to be used in tank mixture with other products, follow the respective manufacturers' label recommendations. Where wettable powder formulations are used in combination with **FARM-AG FOLPET 500 SC**, slurry should firstly be made with the powder and some water, mixed into the spray tank and finally the **FARM-AG FOLPET 500 SC** can be added. Tank mixtures must be sprayed out immediately and not allowed to stand in the spray tank. Spraying equipment must be thoroughly flushed out at the end of the spraying operation.

Application:

- Ensure that the equipment is correctly calibrated and is checked regularly during application to ensure even and accurate application.
- Apply as a preventative application and ensure thorough coverage of both upper and lower leaf surfaces.
- **FARM-AG FOLPET 500 SC** can be applied with conventional high volume spray equipment. It is important that the sprayer be in good working order and correctly calibrated prior to application, to ensure that the correct delivery rate and even distribution of the spray mixture is obtained.
- Constant agitation throughout the spray operation is essential.

CROP AND DISEASE	DOSAGE	REMARKS
Table grapes: Deadarm disease, Anthracnose and Downy mildew	200 mL / 100 L water	Recommended only until pre-blossom stage. Apply first spray at budswel stage. Repeat application at 7 – 10 day intervals depending on weather conditions.

WAARSKUWINGS

Onthoudingsperiodes (Minimum aantal dae tussen laaste toediening en oes of voer/beweiding)

Tafeldruwe	Nie later as voorblomstadium
------------	------------------------------

- Hanteer versigtig.
- Irriterend vir oë en mag sensitisering veroorsaak deur velkontak.
- Giftig vir visse.
- Berg in 'n koel plek en weg van voedsel en voer en nader landboumiddels
- Hou buite bereik van kinders, oningesigte persone en diere.
- Herbetreding: Moenie die behandelde gebied betree totdat die sputnewel droog is tensy beskermende klere gedra word.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormalle klimaats- en bergingstoestande, kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die siekte teen die betrokke middel sowel as die metode, tyd en akkuraathed van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effek op mense of diere of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruikier om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

VOORSORGMAATREËLS

- Vermi kontak met oë envel en inasem van sput newel
- Dra rubberhandskoene en gesigskerm wanneer die konsentraat hanteer word.
- Indien dit in u oë kom, moet dit dadelik uitgespoel word.
- Indien dit op u vel kom, moet dit dadelik afgewas word.
- Indien produk op u klere kom, trek uit en was dadelik.
- Moet nie eet, drink of rook terwyl die produk hanteer word of voordat hande en gesig gewas is en skoon klere aangetrek is nie.
- Voorkom dat die sputnewel orwaai na ander gewasse, weiding, riviere of damme en gebiede wat nie behandel word nie.
- Maak die sputapparaat deeglik skoon na gebruik en gooi waswater waar dit nie voedsel, gewasse, weiding, riviere of damme sal besoedel nie.
- Keer houer om oor die sputtenk om mengtenk om en maak seker die houer is leeg voor dit vernietig word
- **DRIE KEER SPOEL** Keer die leë houer om oor die sput- of mengtenk en dreineer vir ten minste 30 sekondes nadat die vloeistof 'n gedrup verminder het. Spoel die houer daarna drie keer uit met 'n volume water gelijkstaande aan 'n minimum van 10 % van die inhoud van die houer. Gooi die spoelwater by die inhoud van die sputtenk voordat die houer vernietig word op die voorgeskrewe manier
- Vernietig leë houer deur te deurboor en plat te slaan.
- **Moet nooit** die houer vir enige ander doel gebruik nie
- Voorkom kontaminasie van voedsel, voer, drinkwater en eetgerei.

NOODHULP BEHANDELING

OOG KONTAK	Spoel dadelik met ruim hoeveelheid koue water, ook onder oogledle vir ten minste 15 minute, verwyder kontak lense. Kry dadelik mediese hulp.
INAME	Spoel mond uit en drink dan ruim hoeveelheid water. Moenie braaking veroorsaak nie tensy deur gif beheer sentrum of doktor aan bevel word. Kry mediese advies. Moet nooit braaking veroorsaak of enigste persoon vir mond vir bewusteloze persoon, of een wat stuiptrekkings het, gee nie
VEL KONTAK	Trek dadelik besoedelde klere uit. Spoel af met genoegsame water. As vel irritase aanhou kontak mediese hulp. Was besoedelde klere voor hergebruik.
INASEMING	Verwyder persoon na vars lug. Indien asemhaaling ongereeld is of gestopt het, dien kunsmatige asemhaaling toe. Hou patient warm en rustig. Kry mediese hulp of Kontak gif Inligting Sentrum dadelik.

WEERSTANDSWAARSKUWING

FARM-AG FOLPET 500 SC is 'n groepkode M4 swamddoder. Enige populasie van 'n spesifieke onkruid mag individue insluit wat 'n natuurlike weerstand teen **FARM-AG FOLPET 500 SC**, of enige ander groepkode M4 swamddoder het. Indien hierdie groepkode M4 swamddoder herhaaldelik aangewend word, kan die weerstandbiedende individue uiteindelik die swam populasie oorheers. Hierdie weerstandbiedende swamme sal waarskynlik nie deur **FARM-AG FOLPET 500 SC** of enige ander groepkode M4 swamddoder beheer word nie.

Om weerstand teen swamdoders te vertraag:

- Vermy die eksklusiewe herhaalde gebruik van swamdoders met dieselfde groepkode. Wissel af met, of gebruik tenkmengsels van produkte in verskillende swamdodergroepkodes.
- Integreer ander beheermaatreëls (chemies, verbouing, biologies) in swamdoder programme.

Vir spesifieke inligting oor weerstandsbestuur kontak die registrasiehouer van hierdie produk.

VERENIGBAARHEID

- **FARM-AG FOLPET 500 SC** kan in 'n tenkmengsel toegedien word saam met die meeste insek- en swamdoders wat algemeen gebruik word, met uitsondering van Kalk, Kalkswael en Bordeaux Mengsel.

GEBRUIKSAANWYSINGS Gebruik slegs soos aangedui**Verenigbaarheid:**

- **FARM-AG FOLPET 500 SC** kan in 'n tenkmengsel toegedien word saam met die meeste insek- en swamdoders wat algemeen gebruik word, met uitsondering van kalk, kalkswael en Bordeaux Mengsel.
- Moenie meer as 4 toedienings per seisoen toedien nie.
- Moenie **FARM-AG FOLPET 500 SC** saam met enige ander Botrytis middles op tafel druwe toedien nie.

Meng instruksies:

Skud die houer voordat tenk mengsel voorderei word. Maak die sputtenk halfvol en roer. Voeg dan die benodigde hoeveelheid **FARM-AG FOLPET 500 SC** by, terwyl water aanhoudend geroer word. Maak die sputtenk tot die finale volume vol en roer goed. Moenie brak water gebruik nie. Indien die roerde in sputtenk gestaak word moet dit vir ten minste 5 minute weer aangeskakel wees, alvorens weer met toediening voortgegaan word. As tenkmengsels met ander middels gemaak word, moet die betrokke vervaardiger se etiket gevolg word. Waar benabare poerie in kombinasie met **FARM-AG FOLPET 500 SC** in 'n tenkmengsel toegedien word, moet die benabare poerie eers in 'n aparte houer met klien bietjie water gemeng word en dan eerstelike die mengtenk gevoeg word gevolg deur die **FARM-AG FOLPET 500 SC**. Die tenkmengsel moet onmiddellik uitgespuit word en nie toegelaat word om oornag in die sputtenk te staan nie. Die sputtoerusting moet aan die einde van toediening deeglik uitgewas word.

Toediening:

- Verseker die sputtoerusting is korrek gekalibreer en gereeld nagegaan tydens toediening om akkurate toediening te verseker.
- Dien toe as 'n voor kommende behandeling en versekere deeglike bedekking van die bo en onder oppervlaktes van die blare.
- **FARM-AG FOLPET 500 SC** mag met konvensionele hoe volume sputt toegedien word. Dit is belangrik dat die sputtoerusting in goeie werkende toestand is en vooraf gekalibreer is om die regte aflewering en eweredige verspreiding van spuitstof te verseker.
- Konstante roering van die sputtmengsel tydens toediening is noodsaaklik.

GEWAS & SWAMSIEKTE	DOSIS	AANMERKINGS
Tafel druwe: Streepvleksiekte, Antraknose en Donsige skimmel	200 ml / 100 ℥ water	Slegs aanbeveel tot voor-blom stadium. Dien die eerste bespuiting toe by ogieswelstadium en herhaal met 7 tot 10 dae tussenposes afhangende van weerstoestande.



FARM-AG FOLPET 500 SC

Reg No/ Nr L10591 Act No/ Wet Nr 36/1947

A suspension concentrate organic based fungicide for the preventive control of deadarm disease, anthracnose and downy mildew preblossom in table grapes as listed.

'n Suspensie konsentraat organiesgebaseerde swamddoder vir die voorkomende beheer van streepveleksiwkte, antraknose en donsige skimmel voor-blom in tafeldruwe.

FRAC FUNGICIDE GROUP CODE

M4

FRAC SWAMDODER GROEPKODE

ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL:

Folpet (phtalimide)..... 500 g/l..... Folpet (ptalimied)

Manufacture Date:

Vervaardigingsdatum:

Batch No.:

Lotnommer:

NET MASS/ INHOUD

l

**SEE ENCLOSED LEAFLET FOR FULL PARTICULARS
SIEN INGESLOTE VOUBILJET VIR VOLLE BESONDERHEDE**

REGISTRATION HOLDER/REGISTRASIEHOUER:

Cedar Falls Properties 130 (Pty) Ltd.

Co. Reg.No. 2006/031383/07

61 Marshall Drive, Old Mill Industrial Park, Mt Edgecombe, South Africa, 4300 • Tel. (031) 003 3486

Emergency Number: Griffon Poison Information Centre

UN No.: 3082



**CAUTION
VERSIGTIG**



165 x 140 mm

CYAN

MAG

YEL

BLK