

## WARNINGS

**Hazard Statements:** Harmful if swallowed or inhaled. May be harmful in contact with skin. Causes mild skin irritation. Causes serious eye irritation. Suspected of damaging fertility or the unborn child. Very toxic to aquatic life.

**Pre Harvest Interval:** Allow the following number of days to elapse between the final application and harvest of the crop:

Potatoes and youngberries	14 days
Wine grapes and pineapples	21 Days
Table grapes	56 Days

**NOTE:** The recommended withholding period between last application and harvest will meet local residue requirements when the use instructions are strictly adhered to but may not meet export residue requirements. If the treated crop is intended for the export market, consult the relevant exporting authority regarding pre-harvest interval requirements, before the application of this product.

- Store in a cool place.
- Store away from food and feeds.
- Keep out of reach of children, uninformed persons and animals.

**Re-entry interval:** Do not enter treated area until spray deposit has dried unless wearing protective clothing.

**Although this remedy has been extensively tested under various conditions, the registration holder does not warrant that it will be efficacious because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label as well as by method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal for any reason whatsoever or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.**

## PRECAUTIONS

### Precautionary Statements

**General:** If medical advice is needed, have product container or label at hand. Keep out of reach of children.

**Prevention:** Obtain, read and follow all safety instructions before use. Avoid breathing dust, spray. Wash hands, forearms and face thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Wear protective clothing, eye protection, face protection, protective gloves.

**Response:** IF SWALLOWED: Get medical help. IF INHALED: Remove person to fresh air and keep comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. IF ON SKIN: Get medical help. IF exposed or concerned, get medical advice. If skin irritation occurs: Get medical help. Collect spillage.

**Storage:** Store locked up.

**Disposal:** Dispose of contents and container to an approved waste disposal plant.

- Do not eat, drink, or smoke whilst mixing or applying the product, before washing hands and face and changing clothes.
- Prevent contamination of food, feed, drinking water and eating utensils.
- Avoid drift of spray mist onto other crops, pastures, rivers, dams, bore holes and areas not under treatment.
- Clean applicator after use.
- Dispose of wash water where it will not contaminate crops, pastures, rivers, dams and bore holes.
- **Clean the plastic-lined paper bag as follows:** Shake all the pesticide formulation out of the bag into the spray or mixing tank. Turn the bag inside out. Hold the bag over the spray or mixing tank and rinse the bag with running water for thirty seconds. Shake the rinsate off into the tank. Repeat the rinsing twice. Do not bury, burn or donate the bag to any other parties that may use it as a container for foodstuffs. These bags cannot be recycled and must be incinerated by a licensed incineration facility.
- Destroy empty container and do not use for any other purpose

# METASTAR 700 WP

Reg. No. / Nr L10846 Act No. 36 of 1947/ Wet Nr 36 van 1947

**FUNGICIDE**



**READ THE LABEL AND ATTACHED PACKAGED LEAFLET BEFORE USE  
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

**ONLY FOR COMMERCIAL FARM USE.  
NOT FOR USE WITH ANY HANDHELD EQUIPMENT AND/OR IN RESIDENTIAL AREAS AND AERIAL APPLICATION.**

**FRAC FUNGICIDE GROUP CODE**

**4/M3**

**FRAC SWAMDODER GROEPKODE**

A contact and systemic fungicide combination formulated as a wettable powder for the control of various diseases on crops as indicated under directions for use.

'n Benatbare poeier kontak- en sistemiese swamdoder mengsel vir die beheer van verskeie siektes op gewasse soos aangedui onder gebruiksaanwysings.

### ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL:

Metalaxyl (phenyl amide)..... 100 g/kg.....Metalaksil (fenielamied)  
Mancozeb (dithiocarbamate).....600 g/kg.....Mankoseb (ditiokarbamaat)



**WARNING**

### Hazard Statements:

Harmful if swallowed or inhaled.  
May be harmful in contact with skin.  
Causes mild skin irritation.  
Causes serious eye irritation.  
Suspected of damaging fertility or the unborn child.  
Very toxic to aquatic life.

### Precautionary Statements:

Avoid breathing dust, spray.  
IF SWALLOWED: Get medical help.  
IF INHALED: Remove person to fresh air and keep comfortable for breathing.  
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.  
IF ON SKIN: Get medical help.  
If skin irritation occurs: Get medical help.

**In case of poisoning, call the following number:  
Poison Information Helpline (24 hrs) 0861 555 777  
Disaster Management Number: Spill Tech 086 100 0366**

Batch No:

D.O.M:

D.O.E:

Net Contents:

See reverse of packaging for details.

FOR FULL PARTICULARS SEE ATTACHED PAMPHLET.

VIR VOLLEDIGE BESONDERHEDE KYK AANGEHEGTE ETIKET.

UN No. / VR Nr: 3077



REGISTRATION HOLDER/ REGISTRASIEHOUER

Farm Ag International (Pty) Ltd  
Reg. Nr.: Reg. No. 2005/011761/07  
61 Marshall Drive, Old Mill Industrial Park,  
Mount Edgecombe, South Africa, 4300  
Tel: 031 003 3486



Scan for GHS  
and label details

**BARCODE**

100643

## RELEVANT SUBSTANCES

NAME	PRODUCT IDENTIFIER
Mancozeb, active ingredient	CAS-No.: 8018-01-7
Metalaxyl, active ingredient	CAS-No.: 57837-19-1
Sodium salt of dialkyl naphthalene sulphonic acid	CAS-No.: 25638-17-9
Sodium salt of naphthalene sulphonate formaldehyde condensate	CAS-No.: 68425-94-5

## RESISTANCE WARNINGS

For resistance management, **METASTAR 700 WP** is a group code 4 and M3 fungicide. Any fungus may contain individuals naturally resistant to **METASTAR 700 WP** and other group code 4 and M3 fungicides. The resistant individuals can eventually dominate the fungus if these fungicides are used repeatedly. The resistant fungi may not be controlled by **METASTAR 700 WP** or any other group code 4 and M3 fungicide.

To delay fungicide resistance:

- Avoid exclusive, repeated use of fungicides from the same fungicide group code. Alternate or tank mix with registered products from different fungicide group codes.
- For tank mixing or alternating with other products in the spray program, apply suitably registered products from different fungicide group codes. Adhere to all use instructions of the alternate individual product labels.
- Integrate other control methods (chemical, cultural, biological) into disease control program.

**METASTAR 700 WP** contains Metalaxyl, as one component. Metalaxy is a fungicide with specific activity against late blight (*Phytophthora infestans*) on potatoes and downy mildew (*Phytophthora viticola*) on grapevines and youngberries.

Resistant strains can develop which could lead to unacceptable levels of control of these diseases.

### Caution

Since the occurrence of resistant strains cannot be forecast, users are advised to keep treated crops under close observation.

Should disease control be unacceptable, contact your supplier and immediately switch to another suitably registered fungicide of a different group code.

- Incidents of *Plasmopara viticola* (Downy mildew) resistant to Metalaxyl, have been found in various grape producing areas in the South Western Cape.
- Incidents of *Phytophthora infestans* (late blight) resistant to Metalaxyl, have been found in potatoes in the Gamtoos Valley and the George district.

For specific information on resistance management contact your local distributor or the registration holder of this product.

## MODE OF ACTION

**Mancozeb:** Reacts with, and inactivates, the sulphhydryl groups of amino acids and enzymes of fungal cells, resulting in disruption of lipid metabolism, respiration and production of ATP. Broad-spectrum, non-systemic fungicide with protective action.

**Metalaxyl:** Inhibits protein synthesis in fungi, by interference with the synthesis of ribosomal RNA. Systemic fungicide with protective and curative action, taken up by the leaves, stems and roots.

## USE RESTRICTIONS

- **METASTAR 700 WP** does not control early blight on potatoes.
- **METASTAR 700 WP** should not be used in tunnels, glass houses and nurseries.
- **METASTAR 700 WP** is readily translocated upwards in young active growing plants.
- As the plants start aging and reach senescence, penetration and translocation becomes restricted which will reduce the activity of **METASTAR 700 WP**.
- It is therefore important to use a suitable alternate fungicide when plants reach senescence.
- **METASTAR 700 WP** should not be applied more than four times per season on potatoes, youngberries and grapes.
- Continuous use of **METASTAR 700 WP** on certain soils may lead to accelerated bio-degradation of metalaxyl resulting in reduced activity.
- Do not alternate **METASTAR 700 WP** with other fungicides of the same group code (Phenylamides) which have a similar mode of action as **METASTAR 700 WP**.

**DIRECTIONS FOR USE:** Use only as directed.

**General characteristics:**

**METASTAR 700 WP** is a combination of Metalaxyl, a systematic fungicide with preventative and curative action, and Mancozeb, a residual contact fungicide with preventative action. Metalaxyl is absorbed, by the roots, green stems and leaves of plants and translocated upwards. The Metalaxyl component of **METASTAR 700 WP** applied as a foliar application on the surface of the leaves, penetrates rapidly, normally within one hour.

**COMPATIBILITY**

The compatibility of **METASTAR 700 WP** with other products may be influenced by the formulation of the other products involved as well as the quality of the water. It is therefore important that a compatibility test is carried out prior to application or mixing of **METASTAR 700 WP** tank mixtures.

**MIXING INSTRUCTIONS**

Mix the required quantity of **METASTAR 700 WP** in a separate container, with a small volume of water to form a slurry. Half fill the spray tank with clean water and begin agitation in the spray tank. Pour the slurry through a 50 mesh sieve into the spray tank and then top up the tank to the required volume. Ensure thorough agitation is maintained whilst mixing and throughout spray application.

**APPLICATION:****1. Ground Application:**

All spray operations must be performed with suitable equipment that is in good working order and correctly calibrated to give the desired coverage of the target plants for that particular method of application.

**APPLICATION INSTRUCTIONS: Apply Only As Directed**

CROPS	DOSAGE RATE	REMARKS
<b>GRAPES</b> (Table and wine grapes)  <b>Downy mildew</b> ( <i>Plasmopara viticola</i> )	270 g/100ℓ	Firstly, apply a preventative spray program with fungicides from another group code. Apply <b>METASTAR 700 WP</b> immediately an infection period or disease warning occurs. Do not wait until the disease becomes established before applying <b>METASTAR 700 WP</b> . Repeat at 10-14 day intervals. Increase the spray volume progressively as the crop develops up to the pea-berry stage: 250 to 1000 ℓ/ha for wine grapes and 500 to 1200 ℓ/ha for table grapes ensuring complete coverage of the crop. Do not apply after the end of December Do not apply after the pea berry stage on Table grapes.
<b>POTATOES</b> <b>Late blight</b> ( <i>Phytophthora infestans</i> )	<b>Ground Application</b> 2.0 kg/ha in 500 ℓ/ha	Firstly apply a protective program with fungicides from other group codes and then apply <b>METASTAR 700 WP</b> immediately an infection period or disease warning occurs. Do not wait until the disease becomes established before applying <b>METASTAR 700 WP</b> . Repeat at 10 to 14 day intervals  Ground application: Apply in 500 ℓ/ha water
<b>YOUNGBERRIES</b> <b>Downy mildew</b> ( <i>Perenospora rubi</i> )	270 g / 100 ℓ	<b>METASTAR 700 WP</b> must be applied from approximately two weeks after green tip and repeated at 14 day intervals. Do not wait until the disease becomes established before applying <b>METASTAR 700 WP</b> . Apply as a high volume spray to the point of run-off ensuring complete coverage of the plant.
<b>PINEAPPLES</b> <b>Root and base or heart rot</b> ( <i>Phytophthora spp.</i> )	5.0 kg / ha	Immediately after planting: Apply <b>METASTAR 700 WP</b> as an overall broadcast soil drench in 2000 to 3000 ℓ water/ha. Follow up 3 to 6 months later after planting and repeat at 3 months intervals. <b>Established plants:</b> Apply <b>METASTAR 700 WP</b> in 2000 to 3000 ℓ water/ha at 3 months intervals.



## WAARSKUWINGS

**Gevaarstellings:** Skadelik indien ingesluk of ingeasem. Kan skadelik wees in kontak met die vel. Veroorsaak ligte velirritasie. Veroorsaak ernstige oogirritasie. Vermoede van skade aan vrugbaarheid of die ongebore kind. Baie giftig vir waterlewe.

**Voor Oes Tussenpose:** Laat die volgende dae toe tussen laaste toediening en oes van die gewasse soos aangedui.

Aartappels en Youngberries	14 dae
Wyndruiwe en Pyanappels	21 dae
Tafeldruiwe	56 dae

**LET WEL:** Die aanbevole Voor Oes Tussenpose tussen die laaste toediening en oes sal lokale residu perke nakom wanneer die toedienings aanbevelings streng nagevolg word maar kan dalk nie uitvoer residu vlakke nakom nie. As die behandelde gewas vir uitvoer bestem is raadpleeg die betrokke uitvoer gesag aangaande die vereiste residu vlakke alvorens toedienings gemaak word.

- Berg in 'n koel plek.
- Berg weg van voedsel en voer.
- Hou buite bereik van kinders, oningeligte persone en diere.

**Herbetroding:** Moet nie behandelde gebied betree alvorens die spuitneerslag droog is nie, tensy beskermende oorklere gedra word.

**Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle omstandighede doeltreffend sal wees nie, angesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale grond- klimaats- en bergingstoestande, kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie, sowel as die metode, tyd en akkuraatheid van die toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade van enige aard, aan gewasse, plantegroei, die omgewing of vir nadelige effek op mens of diere van 'n gebrek aan prestasie as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.**

## VOORSORGMAARTEËLS

### Voorsorgstellings

**Algemeen:** Indien mediese advies nodig is, hou die produkhouer of etiket byderhand. Hou buite bereik van kinders.

**Voorkoming:** Verkry, lees en volg alle veiligheidsinstruksies voor gebruik. Vermyn die inaseming van stof en spuitmiddels. Was hande, voorarms en gesig deeglik na hantering. Moet nie eet, drink of rook wanneer hierdie produk gebruik word nie. Vermyn vrystelling in die omgewing. Dra beskermende klere, oogbeskerming, gesigbeskerming en beskermende handskoene.

**Reaksie:** INDIEN INGESLUK: Kry mediese hulp. INDIEN INGEASEM: Verwyder persoon na vars lug en hou gemaklik vir asemhaling. INDIEN IN OË: Spoel versigtig met water vir 'n paar minute. Verwyder kontaklense, indien teenwoordig en maklik om te doen. Gaan voort met spoel. INDIEN OP VEL: Kry mediese hulp. INDIEN blootgestel of bekommerd, kry mediese advies. Indien velirritasie voorkom: Kry mediese hulp. Versamel gemorste vloeistof.

**Berging:** Berg toegesluit.

**Weggooi:** Gooi inhoud en houer weg by 'n goedgekeurde afvalverwyderingsaanleg.

- Moet nie eet, drink of rook tydens vermenging of toediening van die produk of voordat die hande en gesig gewas is en die klere verander is.
- Voorkom besoedeling van kos, voer, drinkwater en eetgerei.
- Vermyn wegdrywing van spuitnewel na ander gewasse, weiveld, riviere, damme, boorgate en plekke wat nie behandel moet word nie.
- Maak die toedieningsapparaat skoon na gebruik.
- Gooi uitspoelwater weg waar dit nie gewasse, weiveld, riviere, damme en boorgate kan besoedel nie.
- **Maak die plastiek-gevoerde papiersak soos volg skoon:** Skud al die plaagdoderformulering uit die sak in die spuit- of mengtenk. Draai die sak binneste buite. Hou die sak oor die spuit- of mengtenk en spoel die sak uit met lopende water vir dertig sekondes. Skud die spoelwater in die tenk af. Herhaal die spoel twee keer. Moenie die sak begrawe, verbrand of skenk aan enige ander partye wat dit as 'n houer vir voedsel kan gebruik nie. Hierdie sakke kan nie herwin word nie en moet deur 'n gelsensieerde verbrandingsfasiliteit verbrand word.
- Vernietig lêë houers en moet dit vir geen ander doel gebruik nie.

## RELEVANTE STOWWE

NAAM	PRODUK IDENEFISEERDER
Mancozeb, aktiewe bestanddeel	CAS-Nr.: 8018-01-7
Metalaxyl, aktiewe bestanddeel	CAS-Nr.: 57837-19-1
Natriumsout van dialkielnaftaleensulfoniese hulpmiddel	CAS-Nr.: 25638-17-9
Natriumsout van naftaleensulfonaat formaldehydekondensaas	CAS-Nr.: 68425-94-5

## WAARSKUWING TEEN WEERSTAND

**METASTAR 700 WP** is 'n groepkode 4 en M3 swamdoder. Enige populasie van 'n spesifieke swam mag individue insluit wat 'n natuurlike weerstand teen **METASTAR 700 WP** en enige ander groepkode 4 en M3 swamdoder het. Indien hierdie swamdoders herhaaldelik aangewend word, kan die weerstandbiedende individue uiteindelik die swam populasie oorheers. Hierdie weerstandbiedende swamme sal waarskynlik nie deur **METASTAR 700 WP** of enige ander groepkode 4 en M3 swamdoder beheer word nie.

Om swamdoder weerstand te vertraag.

- Vermoed die eksklusiewe herhaaldelike gebruik van swamdoders met dieselfde groepkode. Wissel af met, of gebruik tenkingsels van produkte in verskillende swamdoder groepkodes.
- Vir tenkingsels of afwissel met ander produkte in die spuitprogram gebruik toepaslike geregistreerde produkte van verskillende swamdoder groepe. Hou streng by die gebruiksaanbeveling van die produk etikete.
- Integreer ander beheermaatreëls (chemies, verbouing, biologies) in swamdoder programme.

**Metastar 700 WP** bevat Metalaksel as een komponent. Metalaksel is swamdoder met spesifieke aksie teen laat roes (*Phytophthora infestans*) op aartappels en Donsige meeldouw (*Phytophthora viticola*) op druiwe en youngbessies. Weerstandbiedende stamme kan ontwikkel wat sal lei tot onaanvaarbare beheer van hierdie siektes.

## Waarskuwing

Aangesien weerstand nie voorspel kan word nie, moet verbruikers die behandelde gewasse noukeurig dophou.

Indien die beheer van die siekte nie na wense is nie, kontak dadelik u verskaffer en verander na 'n ander onverwante toepaslik geregistreerde swamdoder van 'n ander groepkode.

- Gevalle van *Plasmophthora viticola* (Donsige meeldouw) weerstand teen metalaksel kom voor in verskeie druiwe produseerende areas in die Suid Wes Kaap.
- Gevalle van *Plasmophthora infestans* (Laat roes) weerstand teen metalaksel is gevind op aartappels in die Gamtoos Vallei en die George gebied.

Vir spesifieke inligting oor weerstandsbestuur kontak die registrasiehouer van hierdie produk.

## WYSE VAN AKSIE

**Mankoseb:** Reageer met, en inaktiveer, die sulfhidriël-groepe van aminosure en ensieme van swamselle, wat lei tot ontworting van lipiedmetabolisme, respirasie en produksie van ATP. Breëspektrum, nie-sistemiese swamdoder met beskermende werking.

**Metalaxyl:** Inhibeer proteïensintese in swamme, deur inmenging met die sintese van ribosomale RNA. Sistemiese swamdoder met beskermende en genesende werking, opgeneem deur die blare, stingels en wortels.

## GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui.

- **METASTAR 700 WP** beheer nie vroeë roes op aartappels nie.
- **METASTAR 700 WP** moet nie in tunnels, glashuise en kwekerie gebruik word nie.
- Wanneer die plante volwassenheid bereik en begin verouder, neem die absorpsie en translokasie af wat die effektiwiteit van **METASTAR 700 WP** laat afneem.
- Dit is dus belangrik dat 'n ander swamdoder op plante gebruik word sodra plante begin verouder.
- **METASTAR 700 WP** moet nie meer as vier keer per seisoen op aartappels, youngbessies en druiwe toegedien word nie.
- Aanhoudende gebruik van **METASTAR 700 WP** op sekere grond kan aanleiding gee tot versnelde bio-afbreking van metalaksel wat aanleiding gee tot afname in die aktiwiteit.
- Moet nie **METASTAR 700 WP** afwissel met ander fenielamiede met dieselfde tipe aksie as **METASTAR 700 WP**.

### Algemene eienskappe

**METASTAR 700 WP** is 'n kombinasie van Metalaksil, 'n sistematiese swamdoder met voorkomende en kuratiewe aksies, en Mankoseb, 'n residuële swamdoder met voorkomende aksies. Metalaksil word geabsorbeer deur die wortels, groen stingels en blare van die plant en opwaarts getranslokeer. Indien Metalaksil op blare toegedien word, word dit baie vinnig opgeneem, normaalweg binne een uur.

### VERENIGBAARHEID

Die verenigbaarheid van **METASTAR 700 WP** met enige ander produk kan beïnvloed word deur die formulاسie van die produkte betrokke sowel as die kwaliteit van die water. Dit is dus van uiterste belang dat 'n verenigbaarheids toets uitgevoer word voor toediening.

### MENGINSTRUKSIES

Meng die vereiste hoeveelheid van **METASTAR 700 WP** met 'n klein volume water sodat 'n dun room verkry word. Maak die spuitnek half vol met skoon water, en terwyl gemeng word, voeg die dun room deur 'n 50 maas sif by. Vul die tenk aan tot die nodige volume. Roer goed gedurende volmaak en gedurende bespuiting.

### TOEDIENING

#### 1. Grondtoediening:

**METASTAR 700 WP** kan toegedien word deur konvensionele hoëvolume spuittoerusting. Die toerusting moet gekalibreer word voor toediening om te verseker dat die regte dosis gelewer word. Die verspreiding van die spuitmengsel moet egalig oor die teikengebied wees.

#### TOEDIENINGS AANBEVELINGS: DIEN TOE STRENG VOLGENS VOORSKRIFTE

GEWASSE	DOSIS	OPMERKINGS
<b>DRUIWE</b> (Tafel en Wyndruiwe) <b>Donsige skimmel</b> ( <i>Plasmopara viticola</i> )	270 g/100ℓ	Dien chemies onverwante swamdoders toe in 'n voorkomende program en sluit <b>METASTAR 700 WP</b> in sodra 'n infeksieperiode of siektewaarskuwing plaasvind. Moet nie wag tot die siekte toeneem voordat <b>METASTAR 700 WP</b> toegedien word nie. Herhaal elke 10-14 dae. Vermeerder die spuitvolume geleidelik tot met die ertjiekorrelstadium: 250-1000 ℓ/ha vir wyndruiwe en 500-1200 ℓ/ha vir tafeldruiwe en sorg dat die plante goed bedek word. Moet nie later as einde Desember toegedien word nie. Moet nie na ertjiekorrelstadium op tafeldruiwe spuit nie.
<b>AARTAPPELS</b> <b>Laat roes</b> ( <i>Phytophthora infestans</i> )	<b>Grond Toediening</b> 2.0 kg/ha in 500 ℓ/ha	Dien chemies onverwante swamdoders toe in 'n voorkomende program en sluit <b>METASTAR 700 WP</b> in, sodra 'n infeksieperiode of siektewaarskuwing plaasvind. Moet nie wag tot die siekte toeneem voordat <b>METASTAR 700 WP</b> toegedien word nie. Herhaal elke 10-14 dae.  Dien toe in 500 ℓ/ha water (grondtoediening)
<b>YOUNGBESSIES</b> <b>Donsige skimmel</b> ( <i>Perenospora rubi</i> )	270 g /100 ℓ	<b>METASTAR 700 WP</b> toedienings moet ongeveer twee weke na groenpunt begin. Herhaal met 14 dae tussenposes.  Moet nie wag tot die siekte toeneem voordat <b>METASTAR 700 WP</b> toegedien word nie. Dien toe met hoëvolume tot punt van afloop en maak seker dat plante bedek word.
<b>PYNAPPELS</b> <b>Wortel en stamvrot</b> ( <i>Phytophthora spp.</i> )	5.0 kg / ha	Onmiddellik na plant: Dien <b>METASTAR 700 WP</b> as 'n algehele grond deurdrenkings bespuiting toe in 2000 –3000 ℓ water/ha onmiddellik na plant. Volg op 3-6 maande later en daarna elke 3 maande.  <b>Gevestigde plante:</b> Dien <b>METASTAR 700 WP</b> toe in 2000-3000 ℓ water/ha elke 3 maande



# METASTAR 700 WP

Reg. No. / Nr L10846 Act No. 36 of 1947/ Wet Nr 36 van 1947

**FUNGICIDE**



**READ THE LABEL AND ATTACHED PACKAGED LEAFLET BEFORE USE  
KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

**ONLY FOR COMMERCIAL FARM USE.  
NOT FOR USE WITH ANY HANDHELD EQUIPMENT AND/OR IN RESIDENTIAL AREAS AND AERIAL APPLICATION.**

**FRAC FUNGICIDE GROUP CODE**

**4/M3**

**FRAC SWAMDODER GROEPKODE**

A contact and systemic fungicide combination formulated as a wettable powder for the control of various diseases on crops as indicated under directions for use.

'n Benatbare poeier kontak- en sistemiese swamdoder mengsel vir die beheer van verskeie siektes op gewasse soos aangedui onder gebruiksaanwysings.

### ACTIVE INGREDIENT/AKTIEWE BESTANDEEL:

Metalaxyl (phenyl amide)..... 100 g/kg.....Metalaksil (fenielamied)  
Mancozeb (dithiocarbamate).....600 g/kg.....Mankoseb (ditiokarbamaat)



**WARNING**

### Hazard Statements:

Harmful if swallowed or inhaled.  
May be harmful in contact with skin.  
Causes mild skin irritation.  
Causes serious eye irritation.  
Suspected of damaging fertility or the unborn child.  
Very toxic to aquatic life.

### Precautionary Statements:

Avoid breathing dust, spray.  
IF SWALLOWED: Get medical help.  
IF INHALED: Remove person to fresh air and keep comfortable for breathing.  
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.  
IF ON SKIN: Get medical help.  
If skin irritation occurs: Get medical help.

**In case of poisoning, call the following number:  
Poison Information Helpline (24 hrs) 0861 555 777  
Disaster Management Number: Spill Tech 086 100 0366**

Batch No:

D.O.M:

D.O.E:

Net Contents:

See reverse of packaging for details.

FOR FULL PARTICULARS SEE ATTACHED PAMPHLET.  
VIR VOLLEDIGE BESONDERHEDE KYK AANGEHEGTE ETIKET.

UN No. / VR Nr: 3077



REGISTRATION HOLDER/ REGISTRASIEHOUER  
Farm Ag International (Pty) Ltd  
Reg. Nr.: Reg. No. 2005/011761/07  
61 Marshall Drive, Old Mill Industrial Park,  
Mount Edgecombe, South Africa, 4300  
Tel: 031 003 3486



Scan for GHS  
and label details

**BARCODE**

100643

165(H) x 140(W) mm

CYAN

MAG

YEL

BLK

100643